



The Functional Role of Dreaming in Emotional Processes

Serena Scarpelli, Chiara Bartolacci, Aurora D'Atri, Maurizio Gorgoni and Luigi De Gennaro*

Department of Psychology, Sapienza University of Rome, Rome, Italy

OPEN ACCESS

Edited by:

Giulia Mattavelli,
University of Milano-Bicocca, Italy

Reviewed by:

Roumen Kirov,
Institute of Neurobiology (BAS),
Bulgaria
Jean-Baptiste Eichenlaub,
UMR5105 Laboratoire de Psychologie
et NeuroCognition (LPNC), France

*Correspondence:

Luigi De Gennaro
luigi.degennaro@uniroma1.it

Specialty section:

This article was submitted to
Consciousness Research,
a section of the journal
Frontiers in Psychology

Received: 13 December 2018

Accepted: 15 February 2019

Published: 15 March 2019

Citation:

Scarpelli S, Bartolacci C, D'Atri A,
Gorgoni M and De Gennaro L (2019)
The Functional Role of Dreaming in
Emotional Processes.
Front. Psychol. 10:459.
doi: 10.3389/fpsyg.2019.00459

Dream experience (DE) represents a fascinating condition linked to emotional processes and the human inner world. Although the overlap between REM sleep and dreaming has been overcome, several studies point out that emotional and perceptually vivid contents are more frequent when reported upon awakenings from this sleep stage. Actually, it is well-known that REM sleep plays a pivotal role in the processing of salient and emotional waking-life experiences, strongly contributing to the emotional memory consolidation. In this vein, we highlighted that, to some extent, neuroimaging studies showed that the processes that regulate dreaming and emotional salience in sleep mentation share similar neural substrates of those controlling emotions during wakefulness. Furthermore, the research on EEG correlates of the presence/absence of DE and the results on EEG pattern related to the incorporated memories converged to assign a crucial role of REM theta oscillations in emotional re-processing. In particular, the theta activity is involved in memory processes during REM sleep as well as during the waking state, in line with the continuity hypothesis. Also, the gamma activity seems to be related to emotional processes and dream recall as well as to lucid dreams. Interestingly, similar EEG correlates of DE have been found in clinical samples when nightmares or dreams occur. Research on clinical samples revealed that promoting the rehearsal of frightening contents aimed to change them is a promising method to treat nightmares, and that lucid dreams are associated with an attenuation of nightmares. In this view, DE can defuse emotional traumatic memories when the emotional regulation and the fear extinction mechanisms are compromised by traumatic and frightening events. Finally, dreams could represent a sort of simulation of reality, providing the possibility to create a new scenario with emotional mastery elements to cope with dysphoric items included in nightmares. In addition, it could be hypothesized that the insertion of bizarre items besides traumatic memories might be functional to “impoverish” the negative charge of the experiences.

Keywords: dreaming, emotional memories, REM sleep, PTSD, nightmares, narcolepsy, theta oscillations

INTRODUCTION

Dreaming represents a fascinating experience linked to emotional processes so much so to be considered as a key to “access” in the human inner world (e.g., Freud's view; Freud, 1955). In the last decades, dream experience (DE) has been studied under the scientific perspective, defining dream recall as the retrieval of mental sleep elaboration with a different level of complexity and fragmentation, reported after awakenings (Fagioli, 2002).

For years, researchers considered Rapid Eye Movements (REM) sleep the privileged stage to dream (Desseilles et al., 2011) as long as different criteria to collect the reports were introduced (Foulkes, 1962). Actually, DE could be categorized as dream-like or thought-like on the basis of dream contents (Cavallero et al., 1992), including in the first the mentation related to higher emotionality and narrative contents and in the second, to more fragmented elements (Cipolli et al., 2017). Several studies confirmed that dreaming can occur both during REM and NREM sleep (Foulkes, 1962; Rechtschaffen et al., 1963; Monroe et al., 1965; Pivik and Foulkes, 1968; Taub, 1971; Cavallero et al., 1992; Antrobus et al., 1995; Stickgold et al., 2001). In particular, the DEs were obtained in up to 50% of cases upon awakening from NREM sleep, especially from stage 2 (Foulkes, 1962; Pivik and Foulkes, 1968; Nielsen, 2000) and REM sleep suppression pharmacologically-induced did not affect dream recall (Landolt et al., 2001; Oudiette et al., 2012).

Although the exclusive relation between REM sleep and dream has been overcome (Scarpelli et al., 2015a), several studies point out that emotional and perceptually vivid contents are more frequent when reported upon awakenings from REM sleep (Foulkes et al., 1988; Nielsen et al., 1991; Merritt et al., 1994; Hobson et al., 2000; Kahn et al., 2002). More in general, REM sleep plays a pivotal role in the processing of emotional events and several studies showed that the consolidation of emotional memories occurs in this sleep stage (e.g., Lara-Carrasco et al., 2009; Nishida et al., 2009). In addition, experimental deprivation of REM sleep has been demonstrated to compromise the consolidation of emotional stimuli (Cartwright et al., 1975; Wagner et al., 2001; Lara-Carrasco et al., 2009; Spoormaker et al., 2014).

Dream contents including negative emotions (e.g., anxiety and fear) are more frequent than positive ones (Foulkes et al., 1988; Nielsen et al., 1991; Merritt et al., 1994; Fosse et al., 2001), and they are often related to waking-life experiences (Stickgold et al., 2001; Wamsley et al., 2010; Eichenlaub et al., 2017; Vallat et al., 2017). In this regard, the existing empirical evidence highlighted that the neural activation of emotional-limbic (Nir and Tononi, 2010) and reward systems (Perogamvros and Schwartz, 2012) during REM-DE contributes to the offline reprocessing of emotions and associative learning (Perogamvros et al., 2013).

Interestingly, clinical studies have also provided evidence on the potential interaction among DE, sleep alterations and affective disorders (Levin and Nielsen, 2007; Nielsen and Levin, 2007; Schredl, 2011). For instance, mood disturbances frequently appear along with sleep modifications involving REM sleep (Benca et al., 1992, 1997; Walker and van der Helm, 2009). More directly, abundant dreams or nightmares are related to REM sleep abnormalities and psychiatric disorders (Cartwright et al., 2003; Modell et al., 2005; Agargun et al., 2007; Schredl et al., 2009; Sjöström et al., 2009; Marinova et al., 2014; Nakajima et al., 2014).

Taking into account the impossibility to directly access to the DE, most studies in the last decades were focused on the retrieval of dream contents upon awakenings (Scarpelli et al., 2015a). Specifically, several investigations using spontaneous or provoked awakenings in the laboratory aimed to find the electrophysiological (EEG) patterns related to dream recall,

considering it as a form of episodic-declarative memory (Mangiaruga et al., 2018).

In keeping with some earlier reviews (e.g., Revonsuo, 2000; Revonsuo et al., 2015), it could be hypothesized that DE may play a pivotal role in emotional encoding and regulation, nevertheless the specific function and the neural bases of REM-DEs were directly investigated by very few studies (De Gennaro et al., 2011; Eichenlaub et al., 2018; Vallat et al., 2018a).

In light of the above, here we reviewed the main findings about neural bases of dreaming and the link between EEG correlates of DE and emotional processing, underlying that some evidence on sleep disorders characterized by a peculiar DE, especially occurring in REM sleep, may provide a better understanding on the functional role of dreaming and its potential applications in the clinical field.

METHODOLOGICAL NOTE

Studies were identified via PUBMED queries. Key search terms included:

- “Dreaming” and “REM sleep” and “Neuroimaging” in title/abstract (62 articles)
- “Dreaming” and “REM sleep” and “Emotional processing” (22 articles);
- “Dreaming” and “EEG” in title/abstract (686 articles);
- “Dreaming” and “PTSD” in title/abstract (450 articles);
- “Dreaming” and “Emotional Memories” in title/abstract (61 articles);
- “Dreaming” and “Narcolepsy” in title/abstract (128 articles);

We grouped the identified articles in the following categories:

1. Cognitive features, emotional aspects and EEG patterns of mental activity during REM sleep;
2. Dreaming and emotional process: (a) role of REM sleep in processing salient emotional waking life experiences; (b) EEG patterns implicated in REM sleep and emotional dreams; (c) associations among traumatic events and nightmares; (d) role of lucid dreams;
3. Dreaming in narcolepsy.

We focused on the role of emotions in REM sleep, excluding non-English articles. All the articles resulting from using this method and related to our focus were included. Following these criteria, we identified 88 publications which were estimated to be of interest for further examination.

NEURAL BASES OF DREAMING AND EMOTIONAL PROCESSING

A large body of evidence showed that the regions implicated in emotional processes during wakefulness are also responsible for the neurophysiological background of REM sleep that can explain some qualitative features of DE (Maquet et al., 1996; Nir and Tononi, 2010; De Gennaro et al., 2011, 2016; Eichenlaub et al., 2014; Vallat et al., 2018a). In this section, we highlighted the commonalities between the neural bases of REM-DE and emotional processing.

Earlier studies on anatomical correlates of dream alterations by brain lesions studies confirmed the central role of specific cortical and subcortical areas in DE. The main findings point out that two specific cortical systems underlie dreaming. Firstly, a posterior system involves the Temporo-Parieto-Occipital Junction (TPJ) and lesions located in this region alter both waking mental imagery and sleep (Solms, 1997, 2000). The anterior system includes the ventromedial prefrontal cortex (vmPFC) and the white matter surrounding the anterior horns of the lateral ventricles and its damage seems related to dream loss (Solms, 1997, 2000, 2011). It should be noted that many of the afferent and efferent fibers of these two systems are connected with the limbic system and the ventromesial frontal white matter seems to take a part in the interplay between basal forebrain and limbic structures (De Gennaro et al., 2012, 2016).

More recently, human neuroimaging studies have gained a crucial role in dream research. Positron Emission Tomography (PET) and functional Magnetic Resonance Imaging (fMRI) were used especially in the measurement of functional changes in the brain during REM sleep. It has been demonstrated that the distribution of brain activity in REM sleep is not homogenous (Maquet, 2000; Nir and Tononi, 2010). Brain imaging studies found increased regional brain activity in the limbic and paralimbic structures, pontine tegmentum, thalamus and basal forebrain during REM sleep, as compared to wakefulness (Braun et al., 1997, 1998; Nofzinger et al., 1997; Maquet, 2000) and NREM sleep (Braun et al., 1997, 1998; Maquet et al., 2005). In addition, one of the first neuroimaging studies collecting dream report (Maquet et al., 1996) found a bilateral activation of the amygdala in subjects reporting DE upon awakenings from REM sleep, not providing any comparisons with non-recall conditions or awakenings from NREM sleep.

Moreover, remarkable activation of motor and premotor regions has been found in healthy subjects during REM sleep (Braun et al., 1997; Maquet et al., 2000; De Carli et al., 2016), albeit this sleep stage is characterized by muscular atonia thanks to the inhibition of spinal motor neurons by the ponto-bulbar reticular formation (Lai and Siegel, 1999). This seems in line with the observation that individuals affected by REM sleep behavior disorder (RBD) exhibit motor behaviors linked to their DE (Oudiette et al., 2009). Other brain areas are hypoactive during REM sleep compared to the waking state, such as the dorso-lateral PFC (dlPFC), precuneus, orbitofrontal cortex (OFC) and posterior cingulate gyrus (Maquet et al., 1996; Braun et al., 1997; Nofzinger et al., 1997). This evidence could explain the altered executive functions, time perception, and the lack of insight during DE (Desseilles et al., 2011).

Although it is well-known that dreaming is not restricted to REM sleep, some REM sleep features make it a privileged background for DE, especially characterized by high vividness, bizarreness and emotional load (Carr and Solomonova, 2018). In this respect, it should be underlined that most of the regions involved in emotional memory encoding and consolidation (Phelps and LeDoux, 2005; Armony, 2013) are highly activated during REM sleep. In particular, the higher activation of amygdaloid complexes, hippocampal formation and anterior cingulate cortex (ACC) in REM sleep than in wake and/or NREM

sleep may explain the emotional intensity of DE reported upon REM sleep awakenings (Maquet et al., 1996; Braun et al., 1997; Nofzinger et al., 1997; Desseilles et al., 2011; Corsi-Cabrera et al., 2016). Some authors suggested that these structures may be responsible for the reprocessing and consolidation of emotional experiences during REM sleep (Hobson and Pace-Schott, 2002; van der Helm et al., 2011; Deliens et al., 2014).

According to the idea that DE play a role in the emotional processing, more recently awake fMRI measures revealed that subjects experiencing frequent fear during DE have a higher activation of mPFC and reduced activation in the insula, amygdala and midcingulate cortex when faced with aversive stimuli (Sterpenich et al., 2019). The results are consistent with the view that mPFC should exert an inhibitory control on fear expression by reducing amygdala activity (Phelps et al., 2004). A large amount of dream reports were collected at home and the presence of fear in DE has been considered as a trait-like factor of the participants (Sterpenich et al., 2019).

These results are consistent with the continuity hypothesis, namely the possibility that dreams and wakefulness shared similar mechanisms (Schredl, 2003, 2009; Scarpelli et al., 2015a,b; Mangiaruga et al., 2018; Sterpenich et al., 2019).

In fact, neuroimaging data in waking state have provided strong support for a crucial role of the aforementioned regions in emotional processing (Phan et al., 2004). It has been proposed that in the waking brain subcortical regions such as amygdala, nucleus accumbens, locus coeruleus, pulvinar, along with the ACC and OFC are involved in conscious and unconscious emotional processing (Morris et al., 1998; Vuilleumier et al., 2001; Kirov et al., 2012). Specifically, the unconscious perception of emotional stimuli has been related with the functional integrity of subcortical areas that receive an executive cortical feedback from the cortical network (e.g., dlPFC, OFC, and posterior cingulate cortex) only during wakefulness (Tamietto and De Gelder, 2010).

It is well-known that amygdala plays a pivotal role in emotional regulation. Several studies found that this structure is responsible for detecting, generating, and maintaining fear-related emotions (for a review see Phan et al., 2004). Moreover, the amygdala is important in the coordination of adequate responses to threat and danger and it has also been demonstrated that is critical also for response to stimuli that predict positive and negative future outcomes (Paton et al., 2006), allowing the organism to learn more about a stimulus and organize adaptive behavior (Maren, 2011). Specifically, the amygdala, thanks to the connections with the hippocampus, thalamus, mPFC, and ACC, is involved in control of the encoding and retrieval of affective memories and the physical expression of emotions during the waking state (Misane et al., 2005). Also, the hippocampus has an established role in emotional memory encoding and retrieval. Along with the amygdala, the hippocampal formation mediates the processing and execution of fear memories (Phelps, 2004).

Structural brain imaging approaches offer the possibility to focus on ultrastructural, anatomical measures and inter-individual variability associated with DE. This perspective allows researchers to overcome one of the intrinsic limitations of the study of dreams, concerning the difficult to exactly define the time-coupling between the sleep stages and the occurrence of

DE. Neuroanatomical parameters provide a relatively stable measure to account for some trait-like features of DE. In this vein, the correlations between inter-individual differences in quantitative/qualitative features of DE reported by subjects and structural parameters of limbic areas were observed (De Gennaro et al., 2011, 2016). Once again, the regions engaged in emotional encoding and consolidation showed a strong relationship with dream contents, consistently with the studies revealing that the structural characteristics of hippocampus and amygdala are associated with cognitive and emotional processing in waking tasks (Maguire et al., 2000; Bohbot et al., 2007; Iaria et al., 2008). Specifically, microstructural measures were related to the emotional load, bizarreness and vividness of DE reported. For what concerns emotional intensity, a relation was found between lower structural integrity of the left amygdala and decreased emotional load. Moreover, emotionality was linked to a smaller volume of the left hippocampus and larger volume of the right hippocampus. Also, dream contents characterized by high bizarreness were associated with smaller left amygdala, smaller right hippocampus and lower mean diffusivity of the right amygdala.

These observations were partially replicated on patients with Parkinson's Disease (PD), pointing that PD patients did not differ from healthy subjects with respect to sleep and dream characteristics or the neuroanatomical measures and confirming that vividness is related to the amygdalae and also the thickness of the left mPFC. It should be noted that, along with amygdala, mPFC plays a key role in the acquisition, consolidation and retrieval of fear memory and, specifically, modulates fear learning and extinction (Marek et al., 2013).

Moreover, this study on PD focused on the role of the dopaminergic system in DE. It has been found that higher dopamine agonist dosage was associated with qualitatively impoverished DE, as indicated by reduced bizarreness and emotional load levels (De Gennaro et al., 2016). It is worth noting that one of the main regulator of mesolimbic-dopamine network is the mPFC that makes direct and indirect connections to the amygdala and hippocampus (Patton et al., 2013). In other words, the recent findings on PD confirmed the evidence of an essential interplay between the mPFC and mesolimbic-dopamine system for dreaming, as previously showed by data from patients who underwent prefrontal leukotomy that stopped dreaming (Solms, 2011). In fact, in these patients, the ventromedial white matter containing dopaminergic projections to the frontal lobe were disconnected (Bradley et al., 1958). The importance of mPFC in dream processes has been confirmed in another study using PET (Eichenlaub et al., 2014). They revealed that subjects who recall more frequently their DE (i.e., High Recallers—HR) showed higher regional cerebral blood flow (rCBF) in the mPFC during REM sleep and wakefulness than “Low Recallers” (LR) along with higher rCBF in the TPJ during REM, NREM sleep and wakefulness (Eichenlaub et al., 2014). The structural data obtained by MRI from the same sample of Eichenlaub et al. (2014) and from another research (Vallat et al., 2018b) were analyzed, confirming the relationship between brain anatomical structures and dream recall rate (Vallat et al., 2018a). Specifically, the anatomical measures of the mPFC were significantly different

between HR and LR (i.e., increased mPFC white-matter density in HR compared to LR was found). It should be underlined that mPFC is engaged in mind representations and evaluation (for a review, see Legrand and Ruby, 2009), that could have a crucial role in the emotional processing. More in general, it has been hypothesized that mPFC is related to cognitive aspects of emotional processing, such as appraisal/identification of emotion, attention to emotion, awareness and introspection (Phan et al., 2004). Furthermore, it has been demonstrated the involvement of this region in the social emotions processing (Ruby and Decety, 2004). Recently, it has been demonstrated that traumatic experiences and pathological memories are linked to abnormal interactions between hippocampus and mPFC that are responsible to reduced mnemonic ability and decreased emotional control (Maren et al., 2013; Jin and Maren, 2015).

Moreover, it is worth noting that lesion, imaging and stimulation data revealed that the TPJ is related to emotional processing and especially to the “theory of mind,” empathy and social cognition in the waking state (Saxe and Kanwisher, 2003; Young et al., 2010; Santiesteban et al., 2012; Van Overwalle and Vandekerckhove, 2013; Jeurissen et al., 2014). In particular, TPJ contributes to the mentalizing and emotional state inference/attribution (Donaldson et al., 2015; Ye et al., 2015; Biervoye et al., 2016).

Taken together, these findings revealed that, to some extent, the processes that regulate dreaming and emotional salience in DE share similar neural substrates of those controlling emotions during wakefulness. Moreover, in the light of the results by Eichenlaub et al. (2014), one might speculate that the high dream recaller may be particularly interested in their inner world (Ioannides et al., 2009), given that salience seems to have a pivotal role in dream recall, as evaluated in older individuals measuring episodic memory recall (for a review, see Mangiaruga et al., 2018).

ELECTROPHYSIOLOGICAL CORRELATES OF EMOTIONAL MEMORIES DURING REM SLEEP AND WAKEFULNESS

In keeping with the neuroimaging studies exhibiting the relationship among REM sleep, dreaming and processing of emotional experience (De Gennaro et al., 2011, 2016; Eichenlaub et al., 2014; Vallat et al., 2018a), some investigations aimed to identify the sleep EEG correlates of emotional memory consolidation (Levin and Nielsen, 2007; Lara-Carrasco et al., 2009; Nishida et al., 2009; Walker, 2009; Walker and van der Helm, 2009; Spoormaker et al., 2014; Genzel et al., 2015; Hutchinson and Rathore, 2015; Sopp et al., 2017, 2018). Although recent investigations showed that SWS plays a complementary role in this processing (Cairney et al., 2015; Payne et al., 2015; Yordanova et al., 2017), compelling evidence indicates that the potential EEG marker of emotional processes is the frontal theta activity (5–7 Hz) during REM sleep (Nishida et al., 2009; Prehn-Kristensen et al., 2013; Cowdin et al., 2014; Durrant et al., 2015; Seeley et al., 2016). For instance, Nishida et al. (2009) using a nap protocol, demonstrated that the prefrontal theta EEG activity during REM sleep was associated to a remarkable

consolidation benefit for emotional memories rather than neutral memories. Moreover, the extent of emotional consolidation was significantly correlated with the proportion of REM sleep (Nishida et al., 2009). Also, the emotional consolidation of memory has been investigated in healthy and Attention-Deficit Hyperactivity disorder (ADHD) children compared to healthy adults (Prehn-Kristensen et al., 2013). Stimuli from the International Affective Picture System (IAPS) were presented before sleep and after morning awakening in order to test the emotional memory. The authors found that children with and without ADHD showed more slow oscillations (<1 Hz) during SWS and higher theta power during REM sleep. After merging all healthy subjects (children and adults), a correlation between emotional memory and prefrontal oscillations (i.e., theta and slow oscillations) was found. Conversely, ADHD displayed a negative correlation between performance and prefrontal oscillations (Prehn-Kristensen et al., 2013). Interestingly, this study confirmed the pivotal role of the theta band in emotional consolidation, underlying also that this relation is dysfunctional in the clinical group. Actually, this finding seems relevant since ADHD children have a compromised emotional regulation (Martel, 2009) and the marker during sleep of this deficit could be represented by an abnormal theta activity.

More recently, Sopp et al. (2017) showed that REM theta power correlated with enhanced retention of location memory for emotional images, consistent with the idea that REM theta oscillations reflect the reactivation of emotional mnemonic traces.

The role of the REM theta oscillations in emotional memories was also replicated in a recent study that optogenetically silenced the GABAergic neurons in the medial septum causing the inhibition of memory-associated theta rhythm during REM sleep (Boyce et al., 2016). It has been found that silencing these neurons selectively during a REM critical window after learning impaired the performance fear-conditioned contextual memory and the novel object place recognition in the mice (Boyce et al., 2016).

It is worth noting that taken together, these results are in line with the waking EEG studies showing that the theta rhythm substantially correlates with episodic and emotional memory (Klimesch et al., 1996; Klimesch, 1999; Kirov et al., 2009; Uribe et al., 2011). It has been hypothesized that the theta rhythm is directly connected with some areas engaged in memory processing and, in particular, it represents an EEG marker of hippocampal activity (Klimesch, 1996; Cantero et al., 2003; Rutishauser et al., 2010) modulating neuronal changes in hippocampal formation and in neocortical structures (Mitchell et al., 2008). Humans EEG studies confirmed that the frontal theta EEG activity increases during memory tasks (for a review, see Hsieh and Ranganath, 2014) both in the encoding and retrieval stages of memory processing (Klimesch et al., 1997, 2001; Klimesch, 1999; Mölle et al., 2002; Guderian and Düzel, 2005; Addante et al., 2011; Gruber et al., 2013; White et al., 2013). Interestingly, intracranial-EEG (iEEG) studies highlighted that theta oscillations play a pivotal role also in memory formation (Lega et al., 2012). In particular, a higher hippocampal-cortical theta phase coupling was reported in

correspondence of optimal memory performance (Rutishauser et al., 2010). Furthermore, iEEG recordings demonstrated that theta oscillations modulate the interaction between PFC and the medial temporal lobe, confirming that both these areas could be involved in the recall processes (Anderson et al., 2010). More directly, magnetoencephalography (MEG) investigations showed a strong relation between theta activity and mPFC, suggesting that this area could generate theta oscillations (Asada et al., 1999; Ishii et al., 1999; Nishida et al., 2004).

Recently, it has been demonstrated that also the gamma activity during REM sleep represents a potential marker of emotional processing (van der Helm et al., 2011). For instance, an EEG-fMRI study tested the role of REM sleep in the behavioral and brain reactivity to recent waking emotional experiences comparing a group underwent PSG to a waking-group (van der Helm et al., 2011). The results revealed that: a) sleep decreases amygdala reactivity in response to previously encountered emotional experiences (presented by IAPS) as well as behavioral (subjective rating); b) this attenuation of emotional reactivity is related to mPFC connectivity and c) the reduced emotional reactivity is related to attenuated frontal gamma activity during REM sleep (van der Helm et al., 2011). In particular, the gamma activity could represent a marker of the suppression of central adrenergic neurotransmitters in REM sleep (also implicated in arousal and stress) involved in the encoding of emotional salient events along with activation in amygdala-hippocampal system (Walker and van der Helm, 2009; van der Helm et al., 2011). It has been proposed that the adrenergic reduction during REM sleep is essential to emotional regulation because of its role in defusing affective experiences and decreasing emotional intensity (Walker and van der Helm, 2009; van der Helm et al., 2011). For instance, the failure of this reduction during REM sleep has been found in anxiety disturbances associated to the presence of higher gamma activity, contributing to hyperarousal and amygdala reactivity (Etkin and Wager, 2007; Spoormaker and Montgomery, 2008; Walker and van der Helm, 2009). Quite surprisingly, this study did not find any effect on the theta band (van der Helm et al., 2011).

Also, Marshall et al. (2011) found that theta transcranial direct current stimulation (theta-tDCS) applied during REM sleep increases fast beta/gamma EEG oscillations. The theta-tDCS condition was associated with worse mood (i.e., as measured by Positive and Negative Affect Scale negative scores) on the morning after awakening from sleep, compared to sham (Marshall et al., 2011). In line with (van der Helm et al., 2011), this finding points to an indirect link between higher gamma during REM sleep and affective mental states. Moreover, the widespread increase in gamma activity with theta-tDCS provided evidence for the functional coupling between these two rhythms (Marshall et al., 2011).

It should be noted that waking EEG recordings found that theta and gamma oscillations can interact with each other (i.e., cross-frequency coupling) (Canolty et al., 2006; Cohen, 2008; Kramer et al., 2008; Penny et al., 2008; Young and Eggermont, 2009; Tort et al., 2010; Onslow et al., 2011; for a review see Lisman and Jensen, 2013). The theta/gamma coupling has been shown to be functionally relevant for long-term memory

processes (Tort et al., 2009) contributing to the recall of stored information both in rats (Shirvankar et al., 2010) and in humans (Fries et al., 2013). However, these studies revealed that the coupling occurs between the frontal theta and the posterior gamma oscillations (Fries et al., 2013). Little has been known on theta/gamma coupling in human emotional processing. A waking MEG study on emotional expressions found that both the theta and gamma activity occurring in overlapping areas of amygdala, visual cortex and frontal cortex (Luo et al., 2014). Interestingly, the frontal and visual cortical regions showed differences in the temporal profile of activity of these bands: the event-related synchronization peak appeared significantly later in the theta than in the gamma band (Luo et al., 2014). Bearing in mind these observations, it could be hypothesized that the prefrontal gamma could have a different functional role, concerning the reprocessing/depotentiation useful to decrease the emotional salience of recent waking experiences instead of a consolidation/retention of memory traces.

EEG PATTERN OF DREAM EXPERIENCES

Dreaming could reflect the reactivation and consolidation of memories during sleep (Nielsen and Stenstrom, 2005; Wamsley and Stickgold, 2011). It is well-known that after the first peak in incorporation of waking memories into DE (1 day from the events) called “day-residue effect,” there is a second peak following the experience 6–7 days later, i.e., dream-lag effect (Nielsen et al., 2004; Blagrove et al., 2011; Eichenlaub et al., 2019). This delayed incorporation of daytime events has been found to hold for REM sleep dreams but not NREM-DE (Blagrove et al., 2011; van Rijn et al., 2015). Several findings supported that the incorporation of waking-life memories into dream contents represents this process of memory consolidation that allow the integration of the new mnemonic traces in the older memories (Nielsen and Stenstrom, 2005). Although this fascinating perspective has given rise several hypotheses on the functional role of dreaming, very few studies have investigated the EEG correlates of incorporation in REM-DE, while the most of researches pointed to find the EEG pattern predicting the presence/absence of dream recall.

Multi-electrode recordings and quantitative analysis of the EEG signal showed that higher theta activity (5–8 Hz) during REM sleep is related to dream recall both in nocturnal sleep and naps (Marzano et al., 2011; Scarpelli et al., 2015b). In addition, the reduction of the temporo-parietal alpha (8–11 Hz) activity was found before dream recall (Esposito et al., 2004; Marzano et al., 2011). Takeuchi et al. (2003) reported that during NREM sleep onset periods the dream recallers exhibited attenuated alpha and sigma power over the central area (Takeuchi et al., 2003). Nevertheless, they also found higher alpha and sigma activity localized in the central area during REM sleep onset periods in relation to dream recall (Takeuchi et al., 2003). Also, dream recall seems to be associated to increased alpha activity in correspondence of the occipital area, along with a lower frontal alpha power during REM sleep in a multi-nap protocol (Chellappa et al., 2011). It should be

underlined that the alpha activity is one of the EEG bands most influenced by interindividual differences (De Gennaro et al., 2005) and this seems crucial when the experiment included between-subjects measures (Takeuchi et al., 2003; Esposito et al., 2004; Marzano et al., 2011). However, it should be also noted that the alpha, as well as theta band, play a pivotal role in the retrieval of the episodic mnemonic traces during wakefulness as previously mentioned (Klimesch, 1999; Mölle et al., 2002). In the light of above, it has been hypothesized that the neural mechanisms responsible for dream recall resemble those for the encoding and the retrieval of episodic memories during wakefulness (De Gennaro et al., 2012; Scarpelli et al., 2015a).

Once again, in line with the waking EEG studies, it has been reported that also gamma activity has some implications for the recall of DE (Siclari et al., 2017). In fact, Siclari et al. (2017) revealed that the enhancement of rapid frequencies (>25 Hz) during REM and NREM sleep is related to DE when compared with non-recall. In NREM sleep the high-frequency power increased over the parieto-occipital region along with a strong reduction of low-frequency power. Increased high-frequency power related to DE was observed in frontal and temporal areas during REM sleep. Also, it should be mentioned that DEs including “perceiving dimension” correlated with the high-frequency power over the parietal, occipital and temporal areas, while those including thoughts were related to the frontal regions (Siclari et al., 2017).

These patterns are also consistent with the indications obtained by brain lesions and neuroimaging findings in the field of dream research (Solms, 2000; Eichenlaub et al., 2014). It should be noted that a study by transcranial alternating current stimulation (tACS) revealed that the stimulation on the fronto-temporal area in the gamma band during REM sleep is related to lucid dreams (Voss et al., 2014). Albeit this study confirmed the crucial role of the gamma activity for dreaming, it is worth noting that the stimulated areas are different (i.e., fronto-temporal area instead of posterior region).

Interestingly, a very recent combined high-density EEG/fMRI study identified the EEG activity underlining the DE including fear vs. DE without fear (Sterpenich et al., 2019). The results revealed that DE during REM sleep containing fear, as compared with DE without fear, was associated with decreased delta power over the insula and midcingulate cortex (Sterpenich et al., 2019). The same pattern, along with increased gamma power, was found in the right insula during NREM sleep in the case of DE containing fear (Sterpenich et al., 2019). The insula is involved in social-emotion experience and in visceral state (Chang et al., 2013), contributing in the integration of affective information (Shah et al., 2009; Sterpenich et al., 2019). Not surprisingly, also the activation of midcingulate during REM sleep is consistent with previous imaging data that found a critical activation in sensory and motor cortices (De Carli et al., 2016), since that it is a region involving the behavioral/motor responses to danger (Pereira et al., 2010). This activation could reflect an attempt to reactivate threatening situations with emotional and motor reactions that people experienced during wakefulness (Sterpenich et al., 2019).

Some findings provided compelling evidence on the relation between brain activity and dream contents (Dresler et al., 2011; Horikawa et al., 2013; Siclari et al., 2017; Sterpenich et al., 2019) by different approach (e.g., fMRI and machine learning; EEG source modeling; combined fMRI and near-infrared spectroscopy), while -to our knowledge- only few studies were aimed to directly investigated the EEG correlates of dreams containing day-residues, likely because of the difficulty to study the so-called “incorporation.” Dream contents from REM sleep showed a higher probability to include incorporated waking-life experiences (Schredl, 2006; Malinowski and Horton, 2014; Eichenlaub et al., 2018), especially when related to salient emotional events (Voss and Klimke, 2018). Specifically, Eichenlaub et al. (2018) tested the relationship between the theta activity during REM and incorporation of recent waking-life content in DE. They evaluated both emotional intensity and valence of waking-life experiences, assessing the relation with dream report collecting after REM sleep and SWS (Eichenlaub et al., 2018). This study showed that the frontal theta activity in both all and last REM sleep segments correlated to the number of references to waking experiences, identified by the participants into dream recall. Moreover, the reported dream contents contained a high amount of emotional events, irrespective of positive/negative valence (Eichenlaub et al., 2018). No correlations were found for SWS, confirming a strong link between REM sleep and emotional memories. Albeit previous studies showed the pivotal role of the frontal theta power for dream recall in healthy subjects (Marzano et al., 2011; Scarpelli et al., 2015b), this study reported for the first time a direct link between the theta activity during REM sleep and the extent to which dream content is related to recent experiences. In this vein, it could be stated that dreaming reflects to some extent the emotional memory processing.

This perspective was also strengthened by a recent study investigating the activations of facial mimetic musculature during REM sleep as electrophysiological marker of emotional dreams (Rivera-García et al., 2018). The presence of specific facial expression during this stage can be interpreted as a Dream Enacting Behaviors (DEBs), namely the behavioral enactment of the emotional, verbal or motor components of complex dreams (Nielsen et al., 2009; Nir and Tononi, 2010). A growing body of literature suggests that DEBs are more frequent during nightmares or high emotionally intense dreams, providing a background to study the relation between DE and emotional processing (Nielsen et al., 2009; Rivera-García et al., 2018). Rivera-García et al. (2018) analyzed the facial EMG signals along with the standard PSG measures, finding that the activation of corrugator and zygomaticus facial muscles during REM sleep occurs when the incidence of emotional DE is higher (Rivera-García et al., 2018). Besides, this study -examining a wider range of emotions- confirmed that the peak rates in healthy population corresponded to the high negative emotional contents (Rivera-García et al., 2018). This evidence makes the analysis of DEBs a promising approach to study the emotional components of dreaming.

INSIGHTS FROM SLEEP DISORDERS

Until now, we have discussed recent studies highlighting that REM sleep and some electrophysiological features REM-related may play a critical role in the emotional processes.

It should be emphasized that this evidence may afford insights for the translational research since that some dream and/or REM abnormalities represent the core symptoms of psychiatric and sleep disorders (Benca et al., 1992, 1997; Harvey, 2008; Walker and van der Helm, 2009).

The relation among emotions, dreams and sleep disorders appear of special relevance in the context of Post-Traumatic Sleep Disorder (PTSD; Murkar and De Koninck, 2018). PTSD is characterized by hyperreactivity to emotional stimuli and an inability to extinguish traumatic memories (American Psychiatric Association, 2013) and sleep disturbances and recurrent nightmares are recognized as the hallmark of PTSD (Levin et al., 2010; Germain, 2013).

Taking this into account, the interaction of REM-depend learning and PTSD deserves interest (Breslau et al., 2004; Habukawa et al., 2007; Germain et al., 2008; Spoormaker and Montgomery, 2008; Yetkin et al., 2010; Germain, 2013; Pace-Schott et al., 2015). Specifically, PTSD patients report dreams with vivid images and negative emotions associated with the traumatic events (Germain, 2013). It should be considered that the percentage of REM sleep obtained following fear extinction was demonstrated to predict a decrease in autonomic arousal based on skin conductance (Spoormaker et al., 2010) and the disruption of REM sleep impaired extinction (Spoormaker et al., 2012). In this respect, it has been posited that REM sleep may amplify the altered function of the amygdala in PTSD patients, increasing dysphoric dreams (Germain et al., 2008). Moreover, also the idiopathic nightmares originating in early childhood results from dysfunctional hippocampal-amygdala-prefrontal circuit that controls fears memory formation and extinction (Nielsen and Levin, 2007; Marquis et al., 2017). In this view, the so-called “day-residue” and “dream-lag effect” (Nielsen and Stenstrom, 2005) seem relevant for the understanding of sleep-dependent memory consolidation and emotional balance in PTSD (Cipolli et al., 2017). Accordingly to the hypothesis that this phenomenon is an index of the memory consolidation (Nielsen and Stenstrom, 2005), in the context of PTSD, we argue that this process can fail for the emotional memories related to the traumatic events. In fact, dream-lag incorporation occurs more frequently in association to interpersonal problem-solving and are most evident among individuals who report that their DE affects thoughts and feelings during the waking state (Nielsen and Powell, 1992; Kuiken, 2009). Interestingly, Powell et al. (1993) observed that events characterized by high emotional load appear into dream recall for the first 3 days following the event and then again after a delay of from 6 to 7 days. Moreover, (Domhoff, 1996) evidenced that stress-related dream contents may recur over periods of time lasting from days to years. Hence, dreams can represent a reactivation of waking-life experiences with the function of re-processed the conflictual events and their negative emotional tone (Walker and van der Helm, 2009). On the one hand, it can be speculated that REM sleep can provide

the electrophysiological background to allow the fade of the negative emotional charge of autobiographical memories faster than a positive one (Ritchie and Skowronski, 2008). On the other hand, when REM abnormalities occur this functional emotional process fails. In this view, it has been recently demonstrated that negative waking-life experiences incorporated into dreams were characterized by less emotional intensity when reported as dream report (Vallat et al., 2017), accordingly to the idea that dreams foster the emotional regulation reducing the emotional tone of waking-life memories.

Interestingly, a recent study investigated EEG activity in frequent nightmares recallers and controls, observing that higher slow theta (2–5 Hz) activity over frontal and central areas characterized the nightmares group especially during REM sleep compared to controls (Marquis et al., 2017). The results support the idea that REM frontal theta activity also modulates abnormal dreams (i.e., nightmares) in keeping with the evidence that the prefrontal theta activity is related with negatively balanced dreaming during REM sleep (Hebert et al., 2015). Partly, this appears also consistent with the results on EEG correlates of dream recall in healthy subjects (Marzano et al., 2011; Scarpelli et al., 2015b; Eichenlaub et al., 2018). In addition, the prefrontal theta rhythm seems to be related to emotional coping strategy in trauma-exposed subjects that had never developed PTSD, since that these resilient individuals showed higher theta than clinical sample (Cowdin et al., 2014). It is worth noting that DE were not collected in the latter study, not allowing to disentangle if the adaptive role of theta activity is related only to REM sleep *per se* or to some kind of mental sleep activity.

Along with PTSD, also narcolepsy is characterized by a peculiar DE (Schredl, 1998). The study of DE and its relation with emotional processes in narcoleptic patients can deserve attention since that: a) The cataplexy -one of the main narcoleptic symptoms- is generally triggered by intense emotions (De Zambotti et al., 2014); b) The impairment of the limbic system has been repeatedly found in narcoleptic patients (e.g., Mignot et al., 2002; Joo et al., 2005; Hong et al., 2006; Schwartz et al., 2008; Meletti et al., 2015). Narcoleptic patients often report an abundant production of vivid, bizarre and frightening dreams (Schredl, 2008) with a remarkable amount of aggressive themes (Bourguignon, 1976). Investigations on narcoleptic subjects showed higher negatively toned and bizarre dreams along with more terrifying contents than insomnia patients (Mangiaruga et al., 2018). Some authors stated that narcolepsy emphasized the emotional items of their REM dreams as results of the compromised neurobiological system underling the cognitive-emotional functions (Lee et al., 1993). Specifically, it has been found that DE included more anxiety/fear and more bizarre and vivid contents (Fosse, 2000; Fosse et al., 2002).

Also, nightmares seem to occur with a certain frequency in this population (Schredl, 1998; Schredl et al., 2012; Pisko et al., 2014), likely because of daytime stress due to the impairments caused by narcolepsy (Rak et al., 2015). It has been found that the percentage of nightmares in the narcoleptic group was 33% (Pisko et al., 2014), while the prevalence of nightmares in general population was around 5% (Li et al., 2010; Sandman et al., 2013). Conversely, Meaidi et al. (2016) found no differences in

emotional dream contents and in the frequency of nightmares between narcoleptic and healthy subjects. This study showed a greater amount of lucid dreams in narcoleptic than controls (Meaidi et al., 2016). Albeit conflicting results are reported on this issue (Vogel, 1976; Wamsley et al., 2014), according to recent studies (Rak et al., 2015; Meaidi et al., 2016) it could be hypothesized that the increase in lucid dreaming in narcoleptic patients compared to controls can explain the low incidence of nightmares in this clinical group. Namely, lucid dreams may be considered as a sort of coping strategies to deal with the negative emotional contents of DE. In fact, narcoleptic patients who had experience with lucid dreaming felt that dream lucidity provides relief during nightmares (Rak et al., 2015). In other words, dream lucidity led dreamers to affect oneiric contents, also altering their emotional negative charge (Dresler et al., 2011, 2014).

In line with this evidence, some studies suggested that lucid dreams could be useful as intervention for nightmare disorders (Zadra and Pihl, 1997; Spoormaker et al., 2003; Spoormaker and Van den Bout, 2006). Bearing in mind that the methods to induce lucid dreaming often including the promotion of cortical arousal (Stumbrys et al., 2012; Voss et al., 2014), it may be hypothesized that some sleep narcoleptic features such as fragmented sleep, a short sleep latency and sleep onset REM periods, could facilitate lucid dreams (Rak et al., 2015). In fact, the application of the wake-up-back-to-bed technique (i.e., subjects are awakened in the early morning hours and go back to sleep after a period of wakefulness; LaBerge et al., 1994; Stumbrys et al., 2012) has proved to be effective in inducing lucid dreaming in healthy individuals (Erlacher, 2010). It should be noted, the notion that higher cortical desynchronization and fragmented sleep promote dreams is not new in healthy subjects (Koulack and Goodenough, 1976; Scarpelli et al., 2017; Siclari et al., 2017) as well as in nightmare sufferers (Simor et al., 2013) and in narcoleptic patients (Dodet et al., 2015; D'Atri et al., 2019). Specifically, in narcoleptics the successful dream recall upon awakenings was related to higher cortical arousal over parietal regions compared to unsuccessful dream recall, consistently with the findings on healthy individuals (Siclari et al., 2017).

Interestingly, the idea that the awareness of our own dream contents and the possibility of altering them may be beneficial for nightmare sufferers has been proposed by earlier studies (Kellner et al., 1992; Neidhardt et al., 1992; Krakow et al., 2001). Specifically, over 20 years ago the so-called “Imagery rehearsal treatment (IRT)” was introduced (Kellner et al., 1992). IRT is a cognitive-behavioral technique that effectively reduces chronic nightmares within 6–12 weeks of treatment (Marks, 1978; Kellner et al., 1992; Neidhardt et al., 1992; Krakow et al., 1995, 2001, 2002; Forbes et al., 2001; Germain et al., 2004). This method consists of the imaginal rehearsal of a new dream that erased distressing contents, altering the original scenario of nightmare (Kellner et al., 1992). The nightmare sufferers learn to manage the original scenes, creating a less frightened ending and becoming able to include in the new dreams mastery elements, such as the emotional items aimed to change the overall dream affect or emotional reactions to specific dream characters, events or settings (Germain et al., 2004). However, to our knowledge no studies on neurophysiological

correlates of this emotional/traumatic memories re-processing are available.

CONCLUDING REMARKS

Here, we outlined that dreaming during REM sleep may have a pivotal role in the emotional regulation and emotional memory consolidation, accordingly with some previous works (e.g., Cartwright et al., 1998; Desseilles et al., 2011). The current literature does not provide a homogeneous framework on the link between dreaming, emotional processes and neurobiological correlates, albeit remarkable insights from neuroimaging, electrophysiological data and clinical sample led to some final considerations on the functional role of DE that both in healthy and clinical sample serves to affect the inner well-being.

Firstly, we highlighted that emotional regulation and dreaming share similar neurobiological bases suggesting that the amygdala, hippocampus and mPFC operate in a sort of continuum between wakefulness and REM sleep (De Gennaro et al., 2011, 2016; Eichenlaub et al., 2014; Vallat et al., 2018a; Sterpenich et al., 2019).

DE could be considered as a reactivation of waking-life memories and during REM sleep this “re-play” concerns especially highly emotional contents (Cipolli et al., 2017). However, dreaming is a product of an “internal and virtual fabrication” and not always the memory sources are easily identified during wakefulness, so considering DE only as the replay of past experiences may seem reductive (Kirov, 2013; Hobson et al., 2014). Specifically, Hobson et al. (2014) introduced the notion that dreaming plays a pivotal role in enhancing the possibility to create a “virtual world” during sleep that helps to generate more efficient predictions during waking. Hence, dreams not only can contribute to the consolidation of memories with a great emotional load, but it can also represent a mechanism to simulate the real world as a sort of problem-solving based on emotional coping strategies, according to Revonsuo (2000) and Revonsuo et al. (2015).

The studies on dysphoric dreams seem to provide support in favor of this view. In fact, promoting the rehearsal of frightening contents to change them is a promising method to treat nightmares (Kellner et al., 1992). In other words, dreams can defuse emotional traumatic memories when the emotional regulation and the fear extinction mechanism are compromised by traumatic and frightening events, such as in the case of PTSD (Germain et al., 2004), providing the possibility to create a new scenario enriched with mastery elements (Germain et al., 2004). It should be noted that a previous work suggested that the inclusion of bizarre items in the DE may provide a de-contextualization of experiences to promote a better assimilation of salient emotional contents in the existing memories (Horton and Malinowski, 2015). However, it could be hypothesized that the insert of bizarre items beside to traumatic memories might be functional to “impoverish” the negative charge of the experiences.

More recently, from a neurophysiological perspective it has been proposed that the desynchronized neuronal activity in REM sleep, along with the pattern of brain activation (Maquet, 2000)

allows the generation of new associative cell assemblies in highly activated regions such as the hippocampus (Voss and Klimke, 2018). In particular, this model assumes that the bizarreness and vividness dream contents mirror this involuntary associative activity involving the formation of new cell assemblies due to the down-regulated activity in the frontal lobes during REM sleep (Voss and Klimke, 2018).

Beyond these hypotheses, we underlined that the research on EEG correlates of the presence/absence of DE (Marzano et al., 2011; Scarpelli et al., 2015b) and the results on EEG pattern related to the incorporated memories (Eichenlaub et al., 2018) converged to assign a crucial role of REM theta oscillations in this emotional re-processing. In particular, the theta activity is involved in memory processes during REM sleep as well as during waking state (Klimesch et al., 1996; Marzano et al., 2011) in line with the continuity hypothesis (Schredl, 2003, 2009; Scarpelli et al., 2015a,b; Mangiaruga et al., 2018). Also, both posterior and frontal gamma activity seems to be related to emotional processes and dream recall and -interestingly- this band is related to lucid dreams (Voss et al., 2014; Dodet et al., 2015), that are associated with an attenuation of nightmares (Rak et al., 2015). Moreover, the frontal gamma activity seems to have a role in decreasing the amygdala reactivity (van der Helm et al., 2011).

Interestingly, similar EEG correlates of DE have been found in clinical samples when nightmares or dreams occur. Once again, the theta activity (Marquis et al., 2017) and the EEG characterized by high cortical activation (Simor et al., 2013; D’Atri et al., 2019) appear to be crucial in the production and retrieval of DE.

Bearing in mind all of these considerations, some limitations should be mentioned on the hypothesis that REM-DE plays a key role in emotional processing. Firstly, we have highlighted that very little evidence so far is available on a memory function of REM-dreaming *per se* and for this reason it could be challenging to distinguish the role of REM sleep from those of REM-DE in emotional processing. Actually, the role of REM sleep in the mood regulation is somehow controversial since several studies found that this stage has a depressant effect (Palagini et al., 2013). For instance, awakenings from REM sleep are associated to higher scores in the Hamilton Rating Scale for Depression (McNamara et al., 2010), and the increase in REM density (Riemann et al., 1994; Wurts and Edgar, 2000) with higher EEG fast-frequencies (Riemann et al., 1994) have been recognized as the main neurophysiological correlates of depression during sleep. On the one hand, sleep deprivation has been showed beneficial in patients with major depression (Gillin et al., 2001). On the other hand, sleep loss is also related to a compromised consolidation of negative emotional memories (Sterpenich et al., 2007), and this beneficial effect on the mood is linked with emotional lability and inappropriate emotional reactivity in healthy adults under sleep deprivation (Horne, 1988; Dahl, 1996; Gujar et al., 2011). Although disentangling the issue is out of our purpose, undoubtedly, REM sleep alterations are responsible of emotional imbalance and the maintenance of REM sleep promotes adaptive emotional responses during waking life (Walker and van der Helm, 2009). In this view, dreams during REM sleep may help emotional regulation, especially when negative affects are inserted in DE, helping a sort of “exposure”

and desensitization as in the case of the fear extinction (Levin and Nielsen, 2007; Pace-Schott et al., 2012; Menz et al., 2013).

Furthermore, albeit the dream reports from NREM sleep are often “thought-like” (Hobson et al., 2000), some traces of emotions reflecting the individual’s concerns can be observed also during the SWS, as in the case of pavor nocturnus or night terrors (Gottesmann, 2010; Kirov et al., 2012; Castelnovo et al., 2016). NREM as well as REM sleep is involved in the memory replay (Diekelmann and Born, 2010; Oudiette and Paller, 2013), since that both slow oscillations and sleep spindle have been reported in association with memory consolidation (especially declarative) and synaptic plasticity processes (Diekelmann and Born, 2010).

Hence, we do not wish to exclude that also DE during NREM sleep can be “dream-like” and emotional. We emphasize that dream data revealed that affective items are generally higher in REM than in NREM-DE (Hobson et al., 2000) and that the specific physiological background of REM sleep -in continuity with the waking state- may provide a permissive condition for emotional information to be reorganized.

Furthermore, there are important between-subjects and within-subjects variability (Scarpelli et al., 2015a) in recalling dreams. Although the state- and trait-like factors affect the presence/absence of DE, we underlined that the successful DE with high emotional load is observed in the most of cases when clinical samples are analyzed. Along this vein, the DE features may provide some information on the disease process, as in the case of PTSD. Also, taking into account that specific microstructural features have been related to DE, further examination of the EEG pattern in healthy and clinical sample of dreaming may follow two steps: (a) protocols collecting DE in healthy subjects with EEG at high spatial (i.e., high-density EEG) resolution should be carried out in order to replicate the previous findings on the relationship among specific oscillations (gamma; beta; theta; delta) and the dreaming features (e.g., Scarpelli et al., 2015b; Siclari et al., 2017). Within-subjects design, a control of homeostatic and circadian factors as well as technique to better distinguish rhythmic from EEG background signal (e.g., Better Oscillation method; Kaplan et al., 2001) should be used; (b) Studies on EEG correlates of DE in sleep disorders where dreaming is related to the symptoms should be carried out. Also, longitudinal data could be collected to observe the potential changes in the EEG pattern and dream contents.

In particular, the findings discussed point out that both the theta and gamma EEG activities are essential for dreaming and emotional processes. From a translational perspective, it could be interesting to investigate how these frequency bands modulate emotional and traumatic memories incorporated in dreaming. In fact, it could be useful to implement new methods to change the dream content in something of neutral, not disturbing or positive for patients who suffer from nightmares. For example, the tACS techniques allow eliciting lucid dreams through a frontal gamma activity stimulation (Voss et al., 2014). In this perspective, considering lucid dreams as a sort of coping strategies to deal with the negative emotional contents of DE, it could be promising as intervention for nightmare disorder (Zadra and Pihl, 1997; Spoormaker et al., 2003; Spoormaker and Van den Bout, 2006).

Finally, considering that dreaming is an irregular phenomenon and not always observable, it should be considered that a new frontier to investigate more directly the relationship between dreaming and emotions is represented by the DEB (Rivera-García et al., 2018) that can be a viable access to dreams, overcoming the intrinsic difficulty to study this phenomenon (Arnulf, 2012; Alfonsi et al., 2019). More in general, the parasomnias as RBD may be considered a privileged model to study both REM abnormalities and sleep mentation with high emotional load since that these manifestations are often characterized by aggressiveness and negative emotions.

AUTHOR CONTRIBUTIONS

LD and SS made substantial contributions to the conception and design of the work. LD, SS, CB, AD, and MG contributed by drafting the work and revising it critically for important intellectual content, responsible for the final approval of the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

ACKNOWLEDGMENTS

This work was supported by a grant to SS from BIAL Foundation (grant 2016/2017 No. 32/16).

REFERENCES

- Addante, R. J., Watrous, A. J., Yonelinas, A. P., Ekstrom, A. D., and Ranganath, C. (2011). Prestimulus theta activity predicts correct source memory retrieval. *Proc. Natl. Acad. Sci. U.S.A.* 108, 10702–10707. doi: 10.1073/pnas.1014528108
- Agargun, M. Y., Besiroglu, L., Cilli, A. S., Gulec, M., Aydin, A., Inci, R., et al. (2007). Nightmares, suicide attempts, and melancholic features in patients with unipolar major depression. *J. Affect. Disord.* 98, 267–270. doi: 10.1016/j.jad.2006.08.005
- Alfonsi, V., D’Atri, A., Scarpelli, S., Mangiaruga, A., and De Gennaro, L. (2019). Sleep talking: a viable access to mental processes during sleep. *Sleep Med. Rev.* 44, 12–22. doi: 10.1016/j.smrv.2018.12.001
- American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders DMS V*. Washington, DC: American Psychiatric Association. doi: 10.1176/appi.books.9780890425596.744053
- Anderson, K. L., Rajagovindan, R., Ghacibeh, G. A., Meador, K. J., and Ding, M. (2010). Theta oscillations mediate interaction between prefrontal cortex and medial temporal lobe in human memory. *Cerebral Cortex*. 20, 1604–1612. doi: 10.1093/cercor/bhp223
- Antrobus, J., Kondo, T., Reinsel, R., and Fein, G. (1995). Dreaming in the late morning: Summation of REM and diurnal cortical activation. *Consc. Cogn.* 4, 275–299.
- Armony, J. L. (2013). Current emotion research in behavioral neuroscience: the role(s) of the amygdala. *Emot. Rev.* 7, 280–293. doi: 10.1177/1754073912457208

- Arnulf, I. (2012). REM sleep behavior disorder: motor manifestations and pathophysiology. *Move. Disord.* 27, 677–689. doi: 10.1002/mds.24957
- Asada, H., Fukuda, Y., Tsunoda, S., Yamaguchi, M., and Tonoike, M. (1999). Frontal midline theta rhythms reflect alternative activation of prefrontal cortex and anterior cingulate cortex in humans. *Neurosci. Lett.* 274, 29–32. doi: 10.1016/S0304-3940(99)00679-5
- Benca, R. M., Obermeyer, W. H., Thisted, R. A., and Gillin, J. C. (1992). Sleep and psychiatric disorders: a meta-analysis. *Arch. Gen. Psychiatry* 49, 651–668. doi: 10.1001/archpsyc.1992.01820080059010
- Benca, R. M., Okawa, M., Uchiyama, M., Ozaki, S., Nakajima, T., Shibui, K., et al. (1997). Sleep and mood disorders. *Sleep Med. Rev.* 1, 45–56. doi: 10.1016/S1087-0792(97)90005-8
- Biervoye, A., Dricot, L., Ivanou, A., and Samson, D. (2016). Impaired spontaneous belief inference following acquired damage to the left posterior temporoparietal junction. *Soc. Cogn. Affect. Neurosci.* 11, 1513–1520. doi: 10.1093/scan/nsw076
- Blagrove, M., Henley-Einion, J., Barnett, A., Edwards, D., and Seage, C. H. (2011). A replication of the 5–7 day dream-lag effect with comparison of dreams to future events as control for baseline matching. *Conscious. Cogn.* 20, 384–391. doi: 10.1016/j.concog.2010.07.006
- Bohbot, V. D., Lerch, J., Thorndyraft, B., Iaria, G., and Zijdenbos, A. P. (2007). Gray matter differences correlate with spontaneous strategies in a human virtual navigation task. *J. Neurosci.* 27, 10078–10083. doi: 10.1523/JNEUROSCI.1763-07.2007
- Bourguignon, A. (1976). “Narcolepsy and psychoanalysis,” in *Narcolepsy, proceedings of the First International Symposium*, eds. C. Guilleminault, W. C. Dement WC, and P. Passouant (New York, NY: Spectrum Publications), 257–61.
- Boyce, R., Glasgow, S. D., and Williams, S. (2016). Casual evidence for the role of REM sleep theta rhythm in contextual memory consolidation. *Science* 352, 812–816. doi: 10.1126/science.aad5252
- Bradley, K. C., Dax, E. C., and Walsh, K. W. (1958). Modified leucotomy: report of 100 cases. *Med. J. Aust.* 45, 133–138.
- Braun, A. R., Balkin, T. J., Wesensten, N. J., Carson, R. E., Varga, M., Baldwin, P., et al. (1997). Regional cerebral blood flow throughout the sleep-wake cycle: an H₂O-15 positron emission tomography study. *Brain* 120, 1173–1197. doi: 10.1093/brain/120.7.1173
- Braun, A. R., Balkin, T. J., Wesensten, N. J., Gwady, F., Carson, R. E., Varga, M., et al. (1998). Dissociated pattern of activity in visual cortices and their projections during human rapid eye movement sleep. *Science* 279, 91–95. doi: 10.1126/science.279.5347.91
- Breslau, N., Roth, T., Burduvali, E., Kapke, A., Schultz, L., and Roehrs, T. (2004). Sleep in lifetime posttraumatic stress disorder: a community-based polysomnographic study. *Arch. Gen. Psychiatry* 61, 508–516. doi: 10.1001/archpsyc.61.5.508
- Cairney, S. A., Durrant, S. J., Power, R., and Lewis, P. A. (2015). Complementary roles of slow-wave sleep and rapid eye movement sleep in emotional memory consolidation. *Cerebral Cortex* 25, 1565–1575. doi: 10.1093/cercor/bht349
- Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., et al. (2006). High gamma power is phase-locked to theta oscillations in human neocortex. *Science* 313, 1626–1628. doi: 10.1126/science.1128115
- Cantero, J. L., Atienza, M., Stickgold, R., Kahana, M. J., Madsen, J. R., and Kocsis, B. (2003). Sleep-dependent θ oscillations in the human hippocampus and neocortex. *J. Neurosci.* 23, 10897–10903. doi: 10.1523/JNEUROSCI.23-34-10897.2003
- Carr, M., and Solomonova, E. (2018). “Dream recall and content in different stages of sleep and time-of-night effect,” in *Dreams: Biology, Psychology and Culture*, eds K. Valli, R. Hoss, and R. Gongloff (Santa Barbara, CA; Denver, CO: Greenwood Publishing Group), 167–172.
- Cartwright, R., Baehr, E., Kirkby, J., Pandi-Perumal, S. R., and Kabat, J. (2003). REM sleep reduction, mood regulation and remission in untreated depression. *Psychiatry Res.* 121, 159–167. doi: 10.1016/S0165-1781(03)00236-1
- Cartwright, R., Lloyd, S., Butters, L., Weiner, L., McCarthy, L., and Hancock, J. (1975). The effects of REM time on what is recalled. *Psychophysiology* 12, 561–568. doi: 10.1111/j.1469-8986.1975.tb00047.x
- Cartwright, R., Luten, A., Young, M., Mercer, P., and Bears, M. (1998). Role of REM sleep and dream affect in overnight mood regulation: A study of normal volunteers. *Psychiatry Res.* 81, 1–8. doi: 10.1016/S0165-1781(98)00089-4
- Castelnovo, A., Riedner, B. A., Smith, R. F., Tononi, G., Boly, M., and Benca, R. M. (2016). Scalp and source power topography in sleepwalking and sleep terrors: a high-density EEG study. *Sleep* 39, 1815–1825. doi: 10.5665/sleep.6162
- Cavallero, C., Cicogna, P., Natale, V., Occhionero, M., and Zito, A. (1992). Slow wave sleep dreaming. *Sleep* 15, 562–566. doi: 10.1093/sleep/15.6.562
- Chang, L. J., Yarkoni, T., Khaw, M. W., and Sanfey, A. G. (2013). Decoding the role of the insula in human cognition: functional parcellation and large-scale reverse inference. *Cereb. Cortex* 23, 739–749. doi: 10.1093/cercor/bhs065
- Chellappa, S. L., Frey, S., Knoblauch, V., and Cajochen, C. (2011). Cortical activation patterns herald successful dream recall after NREM and REM sleep. *Biol. Psychol.* 87, 251–256. doi: 10.1016/j.biopsycho.2011.03.004
- Cipolli, C., Ferrara, M., De Gennaro, L., and Plazzi, G. (2017). Beyond the neuropsychology of dreaming: insights into the neural basis of dreaming with new techniques of sleep recording and analysis. *Sleep Med. Rev.* 35, 8–20. doi: 10.1016/j.smrv.2016.07.005
- Cohen, M. X. (2008). Assessing transient cross-frequency coupling in EEG data. *J. Neurosci. Methods* 168, 494–499. doi: 10.1016/j.jneumeth.2007.10.012
- Corsi-Cabrera, M., Velasco, F., Río-Portilla, Y., Armony, J. L., Trejo-Martínez, D., Guevara, M. A., et al. (2016). Human amygdala activation during rapid eye movements of rapid eye movement sleep: an intracranial study. *J. Sleep Res.* 25, 576–582. doi: 10.1111/jsr.12415
- Cowdin, N., Kobayashi, I., and Mellman, T. A. (2014). Theta frequency activity during rapid eye movement (REM) sleep is greater in people with resilience versus PTSD. *Exp. Brain Res.* 232, 1479–1485. doi: 10.1007/s00221-014-3857-5
- Dahl, R. E. (1996). The regulation of sleep and arousal: development and psychopathology. *Dev. Psychopathol.* 8, 3–27. doi: 10.1017/S0954579400006945
- D’Atti, A., Scarpelli, S., Schiappa, C., Pizze, F., Vandi, S., Ferrara, M., et al. (2019). Cortical activation during sleep predicts dream experience in narcolepsy. *Ann. Clin. Transl. Neurol.* doi: 10.1002/actn.3.718
- De Carli, F., Proserpio, P., Morrone, E., Sartori, I., Ferrara, M., Gibbs, S. A., et al. (2016). Activation of the motor cortex during phasic rapid eye movement sleep. *Ann. Neurol.* 79, 326–330. doi: 10.1002/ana.24556
- De Gennaro, L., Cipolli, C., Cherubini, A., Assogna, F., Cacciari, C., Marzano, C., et al. (2011). Amygdala and hippocampus volumetry and diffusivity in relation to dreaming. *Hum. Brain Mapp.* 32, 1458–1470. doi: 10.1002/hbm.21120
- De Gennaro, L., Ferrara, M., Vecchio, F., Curcio, G., and Bertini, M. (2005). An electroencephalographic fingerprint of human sleep. *NeuroImage* 26, 114–122. doi: 10.1016/j.neuroimage.2005.01.020
- De Gennaro, L., Lanteri, O., Piras, F., Scarpelli, S., Assogna, F., Ferrara, M., et al. (2016). Dopaminergic system and dream recall: an MRI study in Parkinson’s disease patients. *Hum. Brain Mapp.* 37, 1136–1147. doi: 10.1002/hbm.23095
- De Gennaro, L., Marzano, C., Cipolli, C., and Ferrara, M. (2012). How we remember the stuff that dreams are made of: neurobiological approaches to the brain mechanisms of dream recall. *Behav. Brain Res.* 226, 592–596. doi: 10.1016/j.bbr.2011.10.017
- De Zambotti, M., Pizze, F., Covassin, N., Vandim, S., Cellini, N., Stegagno, L., et al. (2014). Facing emotions in narcolepsy with cataplexy: haemodynamic and behavioural responses during emotional stimulation. *J. Sleep Res.* 23, 432–440. doi: 10.1111/jsr.12133
- Deliens, G., Gilson, M., and Peigneux, P. (2014). Sleep and the processing of emotions. *P. Exp. Brain Res.* 232:1403. doi: 10.1007/s00221-014-3832-1
- Desseilles, M., Dang-Vu, T. T., Sterpenich, V., and Schwartz, S. (2011). Cognitive and emotional processes during dreaming: A neuroimaging view. *Conscious. Cogn.* 20, 998–1008. doi: 10.1016/j.concog.2010.10.005
- Diekelmann, S., and Born, J. (2010). The memory function of sleep. *Nat. Rev. Neurosci.* 11, 114–26. doi: 10.1038/nrn2762
- Dodet, P., Chavez, M., Leu-Semenescu, S., Gormard, J. L., and Arnulf, I. (2015). Lucid dreaming in narcolepsy. *Sleep* 38, 487–497. doi: 10.5665/sleep.4516
- Domhoff, G. W. (1996). *Finding Meaning in Dreams: A Quantitative Approach*. New York, NY: Plenum.
- Donaldson, P. H., Rinehart, N. J., and Enticott, P. G. (2015). Noninvasive stimulation of the temporoparietal junction: a systematic review. *Neurosci. Biobehav. Rev.* 55, 547–572. doi: 10.1016/j.neubiorev.2015.05.017
- Dresler, M., Eibl, L., Fischer, C. F., Spormaker, V. L., Steiger, A., Czisch, M., et al. (2014). Volitional components of consciousness vary across wakefulness, dreaming and lucid dreaming. *Front. Psychol.* 4:987. doi: 10.3389/fpsyg.2013.00987

- Dresler, M., Koch, S., Wehrle, R., Spoormaker, V. I., Holsboer, F., Steiger, A., et al. (2011). Dreamed movement elicits activation in the sensorimotor cortex. *Curr. Biol.* 21, 1833–1837. doi: 10.1016/j.cub.2011.09.029
- Durrant, S. J., Cairney, S. A., McDermott, C., and Lewis, P. A. (2015). Schema-conformant memories are preferentially consolidated during REM sleep. *Neurobiol. Learn. Mem.* 122, 41–50. doi: 10.1016/j.nlm.2015.02.011
- Eichenlaub, J. B., Cash, S. S., and Blagrove, M. (2017). “Daily life experiences in dreams and sleep-dependent memory consolidation” in *Cognitive Neuroscience of Memory Consolidation*, ed. N. Axmacher, B. Rasch (Switzerland: Springer Cham), 161–172.
- Eichenlaub, J. B., Nicolas, A., Daltrozzo, J., Redouté, J., Costes, N., and Ruby, P. (2014). Resting brain activity varies with dream recall frequency between subjects. *Neuropsychopharmacology* 39, 1594–1602. doi: 10.1038/npp.2014.6
- Eichenlaub, J. B., van Rijn, E., Gaskell, M. G., Lewis, P. A., Maby, E., Malinowski, J. E., et al. (2018). Incorporation of recent waking-life experiences in dreams correlates with frontal theta activity in REM sleep. *Soc. Cogn. Affect. Neurosci.* 13, 637–647. doi: 10.1093/scan/nsy041
- Eichenlaub, J. B., van Rijn, E., Phelan, M., Ryder, L., Gaskell, M. G., Lewis, P. A., et al. (2019). The nature of delayed dream incorporation (‘dream-lag effect’): Personally significant events persist, but not major daily activities or concerns. *J. Sleep Res.* 28:e12697. doi: 10.1111/jsr.12697
- Erlacher, D. (2010). “Lucid dreaming. A neural simulation of the real world,” in *Psychiatric Research Trends: Dreams and Geriatric Psychiatry*, ed D. M. Montez (New York, NY: Nova Science Publishers, Inc.), 151–169.
- Esposito, M. J., Nielsen, T. A., and Paquette, T. (2004). Reduced alpha power associated with the recall of mentation from Stage 2 and Stage REM sleep. *Psychophysiology* 41, 288–297. doi: 10.1111/j.1469-8986.00143.x
- Etkin, A., and Wager, T. D. (2007). Functional neuroimaging of anxiety: a metaanalysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 164, 1476–1488. doi: 10.1176/appi.ajp.2007.07030504
- Fagioli, I. (2002). Mental activity during sleep. *Sleep Med. Rev.* 6, 307–320. doi: 10.1053/smr.2001.0214
- Forbes, D., Creamer, M., and Biddle, D. (2001). The validity of the PTSD checklist as a measure of symptomatic change in combat-related PTSD. *Behav. Res. Ther.* 39, 977–986. doi: 10.1016/S0005-7967(00)00084-X
- Fosse, R. (2000). REM mentation in narcoleptics and normals: an empirical test of two neurocognitive theories. *Conscious. Cogn.* 9, 488–509. doi: 10.1006/ccog.2000.0466
- Fosse, R., Stickgold, R., and Hobson, J. A. (2001). Brain-mind states: reciprocal variation in thoughts and hallucinations. *Psychol. Sci.* 12, 30–36. doi: 10.1111/1467-9280.00306
- Fosse, R., Stickgold, R., and Hobson, J. A. (2002). Emotional experience during rapid-eye-movement sleep in narcolepsy. *Sleep* 25, 724–732. doi: 10.1093/sleep/25.7.724
- Foulkes, D. (1962). Dream reports from different stages of sleep. *J. Abnorm. Soc. Psychol.* 65, 14–25. doi: 10.1037/h0040431
- Foulkes, D., Sullivan, B., Kerr, N. H., and Brown, L. (1988). Appropriateness of dream feelings to dreamed situations. *Cogn. Emot.* 2, 29–39. doi: 10.1080/02699938808415227
- Freud, S. (1955). *The Interpretation of Dreams (J. Strachey, Trans.)*. New York, NY: Basic Books.
- Friese, U., Köster, M., Hassler, U., Martens, U., Trujillo-Barreto, N., and Gruber, T. (2013). Successful memory encoding is associated with increased cross-frequency coupling between frontal theta and posterior gamma oscillations in human scalp-recorded EEG. *NeuroImage* 66, 642–647. doi: 10.1016/j.neuroimage.2012.11.002
- Genzel, L., Spoormaker, V. I., Konrad, B. N., and Dresler, M. (2015). The role of rapid eye movement sleep for amygdala-related memory processing. *Neurobiol. Learn. Mem.* 122, 110–121. doi: 10.1016/j.nlm.2015.01.008
- Germain, A. (2013). Sleep disturbances as the hallmark of PTSD: where are we now? *Am. J. Psychiatry* 170, 372–382. doi: 10.1176/appi.ajp.2012.12040432
- Germain, A., Buysse, D. J., and Nofzinger, E. (2008). Sleep-specific mechanisms underlying posttraumatic stress disorder: Integrative review and neurobiological hypotheses. *Sleep Med. Rev.* 12, 185–195. doi: 10.1016/j.smrv.2007.09.003
- Germain, A., Buysse, D. J., Shear, M. K., Fayyad, R., and Austin, C. (2004). Clinical correlates of poor sleep quality in posttraumatic stress disorder. *J. Trauma. Stress* 17, 477–484. doi: 10.1007/s10960-004-5796-6
- Gillin, J. C., Buchsbaum, M., Wu, J., Clark, C., and Bunney Jr, W. (2001). Sleep deprivation as a model experimental antidepressant treatment: findings from functional brain imaging. *Depress. Anxiety* 14, 37–49. doi: 10.1002/da.1045
- Gottesmann, C. (2010). To what extent do neurobiological sleep-waking processes support psychoanalysis? *Int. Rev. Neurobiol.* 92, 233–290. doi: 10.1016/S0074-7742(10)92012-6
- Gruber, M. J., Gelman, B. D., and Ranganath, C. (2013). States of curiosity modulate hippocampus-dependent learning via the dopaminergic circuit. *Neuron* 84, 486–496. doi: 10.1016/j.neuron.2014.08.060
- Guderian, S., and Düzel, E. (2005). Induced theta oscillations mediate large-scale synchrony with mediotemporal areas during recollection in humans. *Hippocampus* 15, 901–912. doi: 10.1002/hipo.20125
- Gujar, N., Yoo, S. S., Hu, P., and Walker, M. P. (2011). Sleep deprivation amplifies reactivity of brain reward networks, biasing the appraisal of positive emotional experiences. *J. Neurosci.* 31, 4466–4474. doi: 10.1523/JNEUROSCI.3220-10.2011
- Habukawa, M., Uchimura, N., Maeda, M., Kotorii, N., and Maeda, H. (2007). Sleep findings in young adult patients with posttraumatic stress disorder. *Biol. Psychiatry* 62, 1179–1182. doi: 10.1016/j.biopsych.2007.01.007
- Harvey, A. G. (2008). Sleep and circadian rhythms in bipolar disorder: seeking synchrony, harmony, and regulation. *Am. J. Psychiatric* 165, 820–829. doi: 10.1176/appi.ajp.2008.08010098
- Hebert, M., Dale, A. L., Brunst-Taiana, C., Campbell, K., and De Koninck, J. (2015). “Spectral EEG correlates of REM sleep dream positive and negative mood: a promising novel approach,” in *29th Annual Meeting of the Associated Professional Sleep Societies* (Seattle, WA).
- Hobson, J. A., Hong, C. C. H., and Friston, K. J. (2014). Virtual reality and consciousness inference in dreaming. *Frontiers in Psychology* 5, 1133.
- Hobson, J. A., and Pace-Schott, E. F. (2002). The cognitive neuroscience of sleep: neuronal system, consciousness and learning. *Nat. Rev. Neurosci.* 3, 679–693. doi: 10.1038/nrn915
- Hobson, J. A., Pace-Schott, E. F., and Stickgold, R. (2000). Dreaming and the brain: toward a cognitive neuroscience of conscious states. *Behav. Brain Sci.* 23, 793–842. doi: 10.1017/S0140525X00003976
- Hong, S. B., Tae, W. S., and Joo, E. Y. (2006). Cerebral perfusion changes during cataplexy in narcolepsy patients. *Neurology* 66, 1747–1749. doi: 10.1212/01.wnl.0000218205.72668.ab
- Horikawa, T., Tamaki, M., Miyawaki, Y., and Kamitani, Y. (2013). Neural decoding of visual imagery during sleep. *Science* 340, 639–642. doi: 10.1126/science.1234330
- Horne, J. (1988). *Why We Sleep: The Functions of Sleep in humans and Other Mammals*. New York, NY: Oxford medical publications.
- Horton, C. L., and Malinowski, J. E. (2015). Autobiographical memory and hyperassociativity in the dreaming brain: Implications for memory consolidation in sleep. *Front. Psychol.* 6:874. doi: 10.3389/fpsyg.2015.00874
- Hsieh, L. T., and Ranganath, C. (2014). Frontal midline theta oscillations during working memory maintenance and episodic encoding and retrieval. *Neuroimage* 85, 721–729. doi: 10.1016/j.neuroimage.2013.08.003
- Hutchinson, I. C., and Rathore, S. (2015). The role of REM sleep theta activity in emotional memory. *Front. Psychol.* 6:1439. doi: 10.3389/fpsyg.2015.01439
- Iaria, G., Comitteri, G., Pastorelli, C., Pizzamiglio, L., Watkins, K. E., and Carota, A. (2008). Neural activity of the anterior insula in emotional processing depends on the individuals’ emotional susceptibility. *Hum. Brain Mapp.* 29, 363–373. doi: 10.1002/hbm.20393
- Ioannides, A. A., Kostopoulos, G. K., Liu, L., and Fenwick, P. B. C. (2009). MEG identifies dorsal medial brain activations during sleep. *Neuroimage* 44, 455–468. doi: 10.1016/j.neuroimage.2008.09.030
- Ishii, R., Shinosaki, K., Ukai, S., Inouye, T., Ishihara, T., Yoshimine, T., et al. (1999). Medial prefrontal cortex generates frontal midline theta rhythm. *Neuroreport* 10, 675–679. doi: 10.1097/00001756-199903170-00003
- Jeurissen, D., Sack, A. T., Roebroek, A., Russ, B. E., and Pascual-Leone, A. (2014). TMS affects moral judgment, showing the role of DLPFC and TPJ in cognitive and emotional processing. *Front. Neurosci.* 8:18. doi: 10.3389/fnins.2014.00018

- Jin, J., and Maren, S. (2015). Prefrontal-hippocampal interactions in memory and emotion. *Front. Syst. Neurosci.* 9:170. doi: 10.3389/fnsys.2015.00170
- Joo, E., Hong, S. B., Tae, W. S., Kim, J. H., Han, S. J., Cho, Y. W., et al. (2005). Cerebral perfusion abnormality in narcolepsy with cataplexy. *Neuroimage* 28, 410–416. doi: 10.1016/j.neuroimage.2005.06.019
- Kahn, D., Combs, A., and Krippner, S. (2002). Dreaming as a function of chaos-like stochastic processes in the self-organizing brain. *Nonlinear Dyn Psychol. Life Sci.* 6, 311–322. doi: 10.1023/A:1019758527338
- Kaplan, A., Röschke, J., Darkhovsky, B., and Fell, J. (2001). Macrostructural EEG characterization based on nonparametric change point segmentation: application to sleep analysis. *J. Neurosci. Methods* 106, 81–90. doi: 10.1016/S0165-0270(01)00331-4
- Kellner, R., Neidhardt, J., Krakow, B., and Pathak, D. (1992). Changes in chronic nightmares after one session of desensitization or rehearsal instructions. *Am. J. Psychiatry* 149, 659–663. doi: 10.1176/ajp.149.5.659
- Kirov, R. (2013). REM sleep and dreaming functions beyond reductionism. *Behav. Brain Sci.* 36, 621–622. doi: 10.1017/S0140525X13001362
- Kirov, R., Brand, S., Kolev, V., and Yordanova, J. (2012). The sleeping brain and the neural basis of emotions. *Behav. Brain Sci.* 35, 155–156. doi: 10.1017/S0140525X11001531
- Kirov, R., Weiss, C., Siebner, H. R., Born, J., and Marshall, L. (2009). Slow oscillation electrical brain stimulation during waking promotes EEG theta activity and memory encoding. *Proc. Natl. Acad. Sci. USA* 106, 15460–15465. doi: 10.1073/pnas.0904438106
- Klimesch, W. (1996). Memory processes, brain oscillations and EEG synchronization. *Int. J. Psychophysiol.* 24, 61–100. doi: 10.1016/S0167-8760(96)00057-8
- Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain Res. Rev.* 29, 169–195. doi: 10.1016/S0165-0173(98)00056-3
- Klimesch, W., Doppelmayr, M., Russegger, H., and Pachinger, T. (1996). Theta band power in the human scalp EEG and the encoding of new information. *Neuroreport* 7, 1235–1240. doi: 10.1097/00001756-199605170-00002
- Klimesch, W., Doppelmayr, M., Schimke, H., and Ripper, B. (1997). Theta synchronization and alpha desynchronization in a memory task. *Psychophysiology* 34, 169–176. doi: 10.1111/j.1469-8986.1997.tb02128.x
- Klimesch, W., Doppelmayr, M., Yonelinas, A., Kroll, N. E. A., Lazzara, M., Röhm, D., et al. (2001). Theta synchronization during episodic retrieval: neural correlates of conscious awareness. *Cogn. Brain Res.* 12, 33–38. doi: 10.1016/S0926-6410(01)00024-6
- Koulack, D., and Goodenough, D. R. (1976). Dream recall and dream recall failure: an arousal-retrieval model. *Psychol. Bull.* 83, 975–984. doi: 10.1037/0033-2909.83.5.975
- Krakow, B., Kellner, R., Pathak, D., and Lambert, L. (1995). Imagery rehearsal treatment for chronic nightmares. *Behav. Res. Ther.* 33, 837–843. doi: 10.1016/0005-7967(95)00009-M
- Krakow, B., Sandoval, D., Schrader, R., Keuhne, B., McBride, L., Yau, C. L., et al. (2001). Treatment of chronic nightmares in adjudicated adolescent girls in a residential facility. *J. Adolesc. Health* 29, 94–100. doi: 10.1016/S1054-139X(00)00195-6
- Krakow, B., Schrader, R., Tandberg, D., Hollifield, M., Koss, M. P., Yau, C. L., et al. (2002). Nightmare frequency in sexual assault survivors with PTSD. *J. Anxiety Disord.* 16, 175–190. doi: 10.1016/S0887-6185(02)00093-2
- Kramer, M. A., Tort, A. B. L., and Kopell, N. J. (2008). Sharp edge artifacts and spurious coupling in EEG frequency comodulation measures. *J. Neurosci. Methods* 170, 352–357. doi: 10.1016/j.jneumeth.2008.01.020
- Kuiken, D. (2009). “Theories of dream function,” in *The Neuroscience of Sleep*, eds R. Stickgold and M. Walker (New York, NY: Elsevier Inc., Academic Press), 295–301.
- LaBerge, S., Phillips, L., and Levitan, L. (1994). An hour of wakefulness before morning naps makes lucidity more likely. *NightLight* 6, 1–5.
- Lai, Y. Y., and Siegel, J. M. (1999). “Muscle Atonia in REM Sleep” in *Rapid Eye Movement Sleep*, eds B. N. Mallick and S. Inoue (London: Narosa), 69–90.
- Landolt, H. P., Raimo, E. B., Schnierow, B. J., Kelsoe, J. R., Rapaport, M. H., and Gillin, J. C. (2001). Sleep and sleep electroencephalogram in depressed patients treated with phenelzine. *Arch. Gen. Psychiatry* 58, 268–276. doi: 10.1001/archpsyc.58.3.268
- Lara-Carrasco, J., Nielsen, T. A., Solomonova, E., Levrier, K., and Popova, A. (2009). Overnight emotional adaptation to negative stimuli is altered by REM sleep deprivation and is correlated with intervening dream emotions. *J. Sleep Res.* 18, 178–187. doi: 10.1111/j.1365-2869.2008.00709.x
- Lee, J. H., Bliwise, D. L., Lebrecht-Bories, E., Guilleminault, C., and Dement, W. C. (1993). Dream disturbed sleep in insomnia and narcolepsy. *J. Nerv. Ment. Dis.* 181, 320–324.
- Lega, B. C., Jacobs, J., and Kahana, M. (2012). Human hippocampal theta oscillations and the formation of episodic memories. *Hippocampus* 22, 748–761. doi: 10.1002/hipo.20937
- Legrand, D., and Ruby, P. (2009). What is self-specific? Theoretical investigation and critical review of neuroimaging results. *Psychol. Rev.* 116, 252–282. doi: 10.1037/a0014172
- Levin, R., Fireman, G., and Nielsen, T. (2010). Disturbed dreaming and emotion dysregulation. *Sleep Med. Clin.* 5, 229–239. doi: 10.1016/j.jsmc.2010.01.006
- Levin, R., and Nielsen, T. A. (2007). Disturbed dreaming, posttraumatic stress disorder, and affect distress: a review and neurocognitive model. *Psych. Bull.* 133, 482–528. doi: 10.1037/0033-2909.133.3.482
- Li, S. X., Zhang, B., Li, A. M., and Wing, Y. K. (2010). Prevalence and correlates of frequent nightmares: a community-based 2-phase study. *Sleep* 33, 774–780. doi: 10.1093/sleep/33.6.774
- Lisman, J. E., and Jensen, O. (2013). The theta-gamma neural code. *Neuron* 77, 1002–1016. doi: 10.1016/j.neuron.2013.03.007
- Luo, Q., Cheng, X., Holroyd, T., Xu, D., Carver, F., and Blair, R. J. (2014). Theta band activity in response to emotional expressions and its relationship with gamma band activity as revealed by MEG and advanced beamformer source imaging. *Front. Hum. Neurosci.* 7: 940. doi: 10.3389/fnhum.2013.00940
- Maguire, E. A., Mummery, C. J., and Büchel, C. (2000). Patterns of hippocampal-cortical interaction dissociate temporal lobe memory subsystems. *Hippocampus* 10, 475–482. doi: 10.1002/1098-1063(2000)10:4<475::AID-HIPO14>3.0.CO;2-X
- Malinowski, J. E., and Horton, C. L. (2014). Evidence for the preferential incorporation of emotional waking-life experiences into dreams. *Dreaming* 24, 18–31. doi: 10.1037/a0036017
- Mangiaruga, A., Scarpelli, S., Bartolacci, C., and De Gennaro, L. (2018). Spotlight on dream recall: the ages of dreams. *Nat. Sci. Sleep* 10, 1–12. doi: 10.2147/NSS.S135762
- Maquet, P. (2000). Functional neuroimaging of normal human sleep by positron emission tomography. *J. Sleep Res.* 9, 207–232. doi: 10.1046/j.1365-2869.2000.00214.x
- Maquet, P., Laureys, S., Peigneux, P., Fuchs, S., Petiau, C., Phillips, C., et al. (2000). Experience-dependent changes in changes in cerebral activation during human REM sleep. *Nat. Neurosci.* 3, 831–836. doi: 10.1038/77744
- Maquet, P., Péters, J. M., Aerts, J., Delfiore, G., Degueldre, C., Luxen, A., et al. (1996). Functional neuroanatomy of human rapid eye movement sleep and dreaming. *Nature* 383, 163–166. doi: 10.1038/383163a0
- Maquet, P., Ruby, P., Maudoux, A., Albouy, G., Sterpenich, V., Dang-Vu, T., et al. (2005). Human cognition during REM sleep and the activity profile within frontal and parietal cortices: a reappraisal of functional neuroimaging data. *Prog. Brain Res.* 150, 219–595. doi: 10.1016/S0079-6123(05)50016-5
- Marek, R., Strobel, C., Bredy, T. W., and Sah, P. (2013). The amygdala and medial prefrontal cortex: partners in the fear circuit. *J. Physiol.* 509, 2381–2391. doi: 10.1113/jphysiol.2012.248575
- Maren, S. (2011). Seeking a spotless mind: extinction, deconsolidation, and erasure of fear memory. *Neuron* 70, 830–845. doi: 10.1016/j.neuron.2011.04.023
- Maren, S., Phan, K. L., and Liberzon, I. (2013). The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nat. Rev. Neurosci.* 14, 417–428. doi: 10.1038/nrn3492
- Marinova, P., Koychev, I., Laleva, L., Kancheva, L., Tsvetkov, M., Bilyukov, R., et al. (2014). Nightmares and suicide: predicting risk in depression. *Psychiatria Danubina* 26, 159–164.
- Marks, I. (1978). Rehearsal relief of a nightmare. *Br. J. Psychiatry* 133, 461–465.
- Marquis, L. P., Paquette, T., Blanchette-Carrière, C., Dumel, G., and Nielsen, T. (2017). REM sleep theta changes in frequent nightmare recallers. *Sleep* 40:zsx110. doi: 10.1093/sleep/zsx110
- Marshall, L., Kirov, R., Brade, J., Mölle, M., and Born, J. (2011). Transcranial electrical currents to probe EEG brain rhythms and

- memory consolidation during sleep in humans. *PLoS ONE*. 6:e16905. doi: 10.1371/journal.pone.0016905
- Martel, M. M. (2009). Research review: A new perspective on attention-deficit/hyperactivity disorder: emotion dysregulation and trait models. *J. Child Psychol. Psychiatry* 50, 1042–1051. doi: 10.1111/j.1469-7610.2009.02105.x
- Marzano, C., Ferrara, M., Mauro, F., Moroni, F., Gorgoni, M., Tempesta, D., et al. (2011). Recalling and forgetting dreams: theta and alpha oscillations during sleep predict subsequent dream recall. *J. Neurosci.* 31, 6674–6683. doi: 10.1523/JNEUROSCI.0412-11.2011
- McNamara, P., Johnson, P., McLaren, D., Harris, E., Beauharnais, C., and Auerbach, S. (2010). REM and NREM sleep mentation. *Int. Rev. Neurobiol.* 92, 69–86. doi: 10.1016/S0074-7742(10)92004-7
- Meadi, A., Pourhadi, N., Ptito, M., Kupers, R., and Jennum, P. (2016). Dreams in narcoleptic patients with and without cataplexy. *J. Sleep Res.* 25, 5–376
- Meletti, S., Vaudano, A. E., Pizzi, F., Ruggieri, A., Vandi, S., Teggi, A., et al. (2015). The brain correlates of laugh and cataplexy in childhood narcolepsy. *J. Neurosci.* 35, 11583–11594. doi: 10.1523/JNEUROSCI.0840-15.2015
- Menz, M. M., Rihm, J. S., Salari, N., Born, J., Kalisch, R., Pape, H. C., et al. (2013). The role of sleep and sleep deprivation in consolidating fear memories. *Neuroimage* 75, 87–96. doi: 10.1016/j.neuroimage.2013.03.001
- Merritt, J. M., Stickgold, R., Pace-Schott, E., Williams, J., and Hobson, J. A. (1994). Emotion Profiles in the Dreams of Men and Women. *Conscious. Cogn.* 3, 46–60. doi: 10.1006/ccog.1994.1004
- Mignot, E., Lammers, G. J., Ripley, B., Okun, M., Nevsimanova, S., Overeem, S., et al. (2002). The role of cerebrospinal fluid hypocretin measurement in the diagnosis of narcolepsy and other hypersomnias. *Arch. Neurol.* 59, 1553–1562. doi: 10.1001/archneur.59.10.1553
- Misane, I., Tovote, P., Meyer, M., Spiess, J., Ögren, S. O., and Stiedl, O. (2005). Time-dependent involvement of the dorsal hippocampus in trace fear conditioning in mice. *Hippocampus* 15, 418–426. doi: 10.1002/hipo.20067
- Mitchell, D. J., McNaughton, N., Flanagan, D., and Kirk, I. J. (2008). Frontal mid-line theta from the perspective of hippocampal theta. *Prog. Neurobiol.* 86, 156–185. doi: 10.1016/j.pneurobio.2008.09.005
- Modell, S., Ising, M., Holsboer, F., and Lauer, C. J. (2005). The Munich vulnerability study on affective disorders: Premorbid polysomnographic profile of affected high-risk probands. *Biol. Psychiatry* 58, 694–699. doi: 10.1016/j.biopsych.2005.05.004
- Möller, M., Marshall, L., Gais, S., and Born, J. (2002). Grouping of spindle activity during slow oscillations in human non-rapid eye movement sleep. *J. Neurosci.* 22, 10941–10947. doi: 10.1523/JNEUROSCI.22-24-10941.2002
- Monroe, L. J., Rechtschaffen, A., Foulkes, D., and Jensen, J. (1965). Discriminability of REM and NREM reports. *J. Personal. Soc. Psychol.* 2, 456–460. doi: 10.1037/h0022218
- Morris, J. S., Öhman, A., and Dolan, R. J. (1998). Conscious and unconscious emotional learning in the human amygdala. *Nature* 393, 467–470.
- Murkar, A. L. A., and De Koninck, J. (2018). Consolidative mechanisms of emotional processing in REM sleep and PTSD. *Sleep Med. Rev.* 41, 173–184. doi: 10.1016/j.smrv.2018.03.001
- Nakajima, S., Mamo, D., Uchida, H., Suzuki, T., Rajji, T., Caravaggio, F., et al. (2014). The optimal dopamine D2 receptor occupancy by antipsychotics in patients with late-life schizophrenia: A [11C]-Raclopride PET study. *Biol. Psychiatry* 75:105S.
- Neidhardt, E. J., Krakow, B., Kellner, R., and Pathak, D. (1992). The beneficial effects of one treatment session and recording of nightmares on chronic nightmare sufferers. *Sleep* 15, 470–473. doi: 10.1093/sleep/15.5.470
- Nielsen, T., and Levin, R. (2007). Nightmares: a new neurocognitive model. *Sleep Med. Rev.* 11, 295–310. doi: 10.1016/j.smrv.2007.03.004
- Nielsen, T., Svob, C., and Kuiken, D. (2009). Dream-enacting behaviors in a normal population. *Sleep* 32, 1629–1636. doi: 10.1093/sleep/32.12.1629
- Nielsen, T. A. (2000). A review of mentation in REM and NREM sleep: “covert” REM sleep as a possible reconciliation of two opposing models. *Behav. Brain Sci.* 23, 851–866. doi: 10.1017/S0140525X0000399X
- Nielsen, T. A., Deslauriers, D., and Baylor, G. W. (1991). Emotions in dream and waking event reports. *Dreaming* 1:287. doi: 10.1037/h0094340
- Nielsen, T. A., Kuiken, D., Alain, G., Stenstrom, P., and Powell, R. A. (2004). Immediate and delayed incorporations of events into dreams: further replication and implications for dream function. *J. Sleep Res.* 13, 327–336. doi: 10.1111/j.1365-2869.2004.00421.x
- Nielsen, T. A., and Powell, R. A. (1992). The day-residue and dream-lag effects: a literature review and limited replication of two temporal effects in dream formation. *Dreaming* 2:67. doi: 10.1037/h0094348
- Nielsen, T. A., and Stenstrom, P. (2005). What are the memory sources of dreaming? *Nature* 437:1286. doi: 10.1038/nature04288
- Nir, Y., and Tononi, G. (2010). Dreaming and the brain: from phenomenology to neurophysiology. *Trends Cogn. Sci.* 14, 88–100. doi: 10.1016/j.tics.2009.12.001
- Nishida, M., Hirai, N., Miwakeichi, F., Maehara, T., Kawai, K., Shimizu, H., et al. (2004). Theta oscillation in the human anterior cingulate cortex during all-night sleep: an electrocorticographic study. *Neurosci. Res.* 50, 331–341. doi: 10.1016/j.neures.2004.08.004
- Nishida, M., Pearsall, J., Buckner, R. L., and Walker, M. P. (2009). REM sleep, prefrontal theta, and the consolidation of human emotional memory. *Cereb. Cortex*. 19, 1158–1166. doi: 10.1093/cercor/bhn155
- Nofzinger, E. A., Mintun, M. A., Wiseman, M., Kupfer, D. J., and Moore, R. Y. (1997). Forebrain activation in REM sleep: an FDG PET study. *Brain Res.* 770, 192–201. doi: 10.1016/S0006-8993(97)00807-X
- Onslow, A. C. E., Bogacz, R., and Jones, M. W. (2011). Quantifying phase-amplitude coupling in neuronal network oscillations. *Prog. Biophys. Mol. Biol.* 105, 49–57. doi: 10.1016/j.pbiomolbio.2010.09.007
- Oudiette, D., De Cock, V. C., Lavault, S., Leu, S., Vidailhet, M., and Arnulf, I. (2009). Nonviolent elaborate behaviors may also occur in REM sleep behavior disorder. *Neurology*. 72, 551–557. doi: 10.1212/01.wnl.0000341936.78678.3a
- Oudiette, D., Dealberto, M. J., Uguccioni, G., Golmard, J. L., Merino-Andreu, M., Tafti, M., et al. (2012). Dreaming without REM sleep. *Conscious. Cogn.* 21, 1129–1140. doi: 10.1002/mds.24044
- Oudiette, D., and Paller, K. A. (2013). Upgrading the sleeping brain with targeted memory reactivation. *Trends Cogn. Sci.* 17, 142–149. doi: 10.1016/j.tics.2013.01.006
- Pace-Schott, E. F., Germain, A., and Milad, M. R. (2015). Sleep and REM sleep disturbance in the pathophysiology of PTSD: The role of extinction memory. *Biol. Mood Anxiety Disord.* 5:3. doi: 10.1186/s13587-015-0018-9
- Pace-Schott, E. F., Verga, P. W., Bennett, T. S., and Spencer, R. M. (2012). Sleep promotes consolidation and generalization of extinction learning in simulated exposure therapy for spider fear. *J. Psychiatr. Res.* 46, 1036–1044. doi: 10.1016/j.jpsychires.2012.04.015
- Palagini, L., Baglioni, C., Ciapparelli, A., Gemignani, A., and Riemann, D. (2013). REM sleep dysregulation in depression: state of the art. *Sleep Med. Rev.* 17, 377–390. doi: 10.1016/j.smrv.2012.11.001
- Paton, J. J., Belova, M. A., Morrison, S. E., and Salzman, C. D. (2006). The primate amygdala represents the positive and negative value of visual stimuli during learning. *Nature* 439, 865–870. doi: 10.1038/nature04490
- Patton, M. H., Bizup, B. T., and Grace, A. A. (2013). The infralimbic cortex bidirectionally modulates mesolimbic dopamine neuron activity via distinct neural pathways. *J. Neurosci.* 33, 16865–16873. doi: 10.1523/JNEUROSCI.2449-13.2013
- Payne, J. D., Kensinger, E. A., Wamsley, E. J., Nathan Spreng, R., Alger, S. E., Gbiler, K., et al. (2015). Napping and the selective consolidation of negative aspects of scenes. *Emotion* 15, 176–186. doi: 10.1037/a0038683
- Penny, W. D., Duzel, E., Miller, K. J., and Ojemann, J. G. (2008). Testing for nested oscillation. *J. Neurosci. Methods* 174, 50–61. doi: 10.1016/j.jneumeth.2008.06.035
- Pereira, M. G., de Oliveira, L., Erthal, F. S., Joffily, M., Mocaiber, I. F., Volchan, E., et al. (2010). Emotion affects action: midcingulate cortex as a pivotal node of interaction between negative emotion and motor signals. *Cogn. Affect. Behav. Neurosci.* 10, 94–106. doi: 10.3758/CABN.10.1.94
- Perogamvros, L., Dang-Vu, T. T., Deseilles, M., and Schwartz, S. (2013). Sleep and dreaming are for important matters. *Front. Psychol.* 4:474. doi: 10.3389/fpsyg.2013.00474
- Perogamvros, L., and Schwartz, S. (2012). The roles of the reward system in sleep and dreaming. *Neurosci. Biobehav. Rev.* 36, 1934–1951. doi: 10.1016/j.neubiorev.2012.05.010
- Phan, K. L., Wager, T. D., Taylor, S. F., and Liberzon, I. (2004). Functional neuroimaging studies of human emotions. *CNS Spectr.* 9, 258–266. doi: 10.1017/S1092852900009196
- Phelps, E. A. (2004). Human emotion and memory: interactions of the amygdala and hippocampal complex. *Curr. Opin. Neurobiol.* 14, 198–202. doi: 10.1016/j.conb.2004.03.015

- Phelps, E. A., Delgado, M. R., Nearing, K. I., and LeDoux, J. E. (2004). Extinction learning in humans: role of the amygdala and vmPFC. *Neuron* 43, 897–905. doi: 10.1016/j.neuron.2004.08.042
- Phelps, E. A., and LeDoux, J. E. (2005). Contributions of the amygdala to emotion processing: from animal models to human behavior. *Neuron* 48, 175–187. doi: 10.1016/j.neuron.2005.09.025
- Piskov, J., Pastorek, L., Buskova, J., Sonka, K., and Nevšimalova, S. (2014). Nightmares in narcolepsy: under investigated symptom? *Sleep Med.* 15, 967–972. doi: 10.1016/j.sleep.2014.03.006
- Pivik, T., and Foulkes, D. (1968). NREM mentation: relation to personality, orientation time, and time of night. *J. Consul. Clin. Psychol.* 32, 144–151. doi: 10.1037/h0025489
- Powell, R. A., Nielsen, T. A., and Cheung, J. (1993). Temporal variations in dream incorporation after a stressful film. *Sleep Res.* 22:139.
- Prehn-Kristensen, A., Munz, M., Molzow, I., Wilhelm, I., Wiesner, C. D., and Baving, L. (2013). Sleep promotes consolidation of emotional memory in healthy children but not in children with attention-deficit hyperactivity disorder. *PLoS ONE* 8:e65098. doi: 10.1371/journal.pone.0065098
- Rak, M., Beiting, P., Steiger, A., Schredl, M., and Dresler, M. (2015). Increased lucid dreaming frequency in narcolepsy. *Sleep* 38, 787–792. doi: 10.5665/sleep.4676
- Rechtschaffen, A., Verdone, P., and Wheaton, J. V. (1963). Reports of mental activity during sleep. *Can. Psychiatr. Assoc. J.* 8, 409–414. doi: 10.1177/070674376300800612
- Revonsuo, A. (2000). The reinterpretation of dreams: An evolutionary hypothesis of the function of dreaming. *Behav. Brain Sci.* 23, 877–901. doi: 10.1017/S0140525X00004015
- Revonsuo, A., Tuominen, J., and Valli, K. (2015). “The Avatars in the Machine - Dreaming as a Simulation of Social Reality,” in *Open MIND*, eds. T. Metzinger and J. M. Windt (Frankfurt am Main: MIND Group).
- Riemann, D., Hohagen, F., Bahro, M., and Berger, M. (1994). Sleep in depression: the influence of age, gender and diagnostic subtype on baseline sleep and the cholinergic REM induction test with RS 86. *Eur. Arch. Psychiatry Clin. Neurosci.* 243, 279–290. doi: 10.1007/BF02191586
- Ritchie, T. D., and Skowronski, J. J. (2008). Perceived change in the affect associated with dreams: the fading affect bias and its moderators. *Dreaming* 18, 27–43. doi: 10.1037/1053-0797.18.1.27
- Rivera-García, A. P., López Ruiz, I. E., Ramírez-Salado, I., González Olvera, J., Guerrero, F. A., and Jiménez-Anguiano, A. (2018). Emotional facial expressions during REM sleep dreams. *J. Sleep Res.* 28:e12716. doi: 10.1111/jsr.12716
- Ruby, P., and Decety, J. (2004). How would You feel versus how do you think She would feel? A neuroimaging study of perspective-taking with social emotions. *J. Cogn. Neurosci.* 16, 988–999. doi: 10.1162/0898929041502661
- Rutishauser, U., Ross, I. B., Mamelak, A. N., and Schuman, E. M. (2010). Human memory strength is predicted by theta-frequency phase-locking of single neurons. *Nature* 464, 903–907. doi: 10.1038/nature08860
- Sandman, N., Valli, K., Kronholm, E., Ollila, H. M., Revonsuo, A., Laatikainen, T., et al. (2013). Nightmares: prevalence among the Finnish general adult population and war veterans during 1972–2007. *Sleep* 36, 1041–1050. doi: 10.5665/sleep.2806
- Santesteban, I., White, S., Cook, J., Gilbert, S. J., Heyes, C., and Bird, G. (2012). Training social cognition: from imitation to theory of mind. *Cognition* 122, 228–235. doi: 10.1016/j.cub.2012.10.018
- Saxe, R., and Kanwisher, N. (2003). People thinking about thinking people: the role of the temporo-parietal junction in “theory of mind”. *Neuroimage* 19, 1835–1842. doi: 10.1016/S1053-8119(03)00230-1
- Scarpelli, S., D’Atri, A., Gorgoni, M., Ferrara, M., and De Gennaro, L. (2015a). EEG oscillations during sleep and dream recall: state- or trait-like individual differences? *Front. Psychol.* 6:605. doi: 10.3389/fpsyg.2015.00605
- Scarpelli, S., D’Atri, A., Mangiaruga, A., Marzano, C., Gorgoni, M., Schiappa, C., et al. (2017). Predicting dream recall: EEG activation during NREM sleep or shared mechanisms with wakefulness? *Brain Topogr.* 30, 629–638. doi: 10.1007/s10548-017-0563-1
- Scarpelli, S., Marzano, C., D’Atri, A., Gorgoni, M., Ferrara, M., and De Gennaro, L. (2015b). State- or trait-like individual differences in dream recall: preliminary findings from a within-subjects study of multiple nap REM sleep awakenings. *Front. Psychol.* 6:928. doi: 10.3389/fpsyg.2015.00928
- Schredl, M. (1998). Dream content in patients with narcolepsy: preliminary findings. *Dreaming* 8, 103–107. doi: 10.1023/B:DREM.0000005900.63728.a7
- Schredl, M. (2003). Continuity between waking and dreaming: a proposal for a mathematical model. *Sleep Hypnos.* 5, 26–39.
- Schredl, M. (2006). Factors affecting the continuity between waking and dreaming: Emotional intensity and emotional tone of the waking-life event. *Sleep Hypnos.* 8, 1–5.
- Schredl, M. (2008). Laboratory references in dreams: Methodological problem and/or evidence for the continuity hypothesis of dreaming? *Int. J. Dream Res.* 1. doi: 10.11588/heidok.00008425
- Schredl, M. (2009). Dreams in patients with sleep disorders. *Sleep Med. Rev.* 13, 215–221. doi: 10.1016/j.smrv.2008.06.002
- Schredl, M. (2011). Dream research in schizophrenia: methodological issues and a dimensional approach. *Conscious. Cogn.* 20, 1036–1041. doi: 10.1016/j.concog.2010.05.004
- Schredl, M., Berger, M., and Riemann, D. (2009). The effect of trimipramine on dream recall and dream emotions in depressive outpatients. *Psychiatry Res.* 167, 279–286. doi: 10.1016/j.psychres.2008.03.002
- Schredl, M., Binder, R., Feldmann, S., Göder, R., Hoppe, J., Schmitt, J., et al. (2012). Dreaming in patients with sleep disorders: a multicenter study. *Somnologie* 16, 32–42. doi: 10.1007/s11818-012-0552-2
- Schwartz, S., Ponz, A., Poryazova, R., Werth, E., Boesiger, P., Khatami, R., et al. (2008). Abnormal activity in hypothalamus and amygdala during humour processing in human narcolepsy with cataplexy. *Brain* 131, 514–522. doi: 10.1093/brain/awn292
- Seeley, W. W., Allman, J. M., Carlin, D. A., Crawford, R. K., Macedo, M. N., Greicius, M. D., et al. (2016). Divergent social functioning in behavioral variant frontotemporal dementia and alzheimer disease: reciprocal networks and neuronal evolution. *Alzheimer Dis. Assoc. Disord.* 21, S50–S57. doi: 10.1097/WAD.0b013e31815c0f14
- Shah, S. G., Klumpp, H., Angstadt, M., Nathan, P. J., and Phan, K. L. (2009). Amygdala and insula response to emotional images in patients with generalized social anxiety disorder. *J. Psychiatry Neurosci.* 34, 296–302.
- Shirvinkar, P. R., Rapp, P. R., and Shapiro, M. L. (2010). Bidirectional changes to hippocampal theta-gamma comodulation predict memory for recent spatial episodes. *Proc. Natl. Acad. Sci. USA.* 107, 7054–7059. doi: 10.1073/pnas.0911184107
- Siclari, F., Baird, B., Perogamvros, L., Bernardi, G., LaRocque, J. J., Riedner, B., et al. (2017). The neural correlates of dreaming. *Nat. Neurosci.* 20, 872–878. doi: 10.1038/nn.4545
- Simor, P., Bódizs, R., Horváth, K., and Ferri, R. (2013). Disturbed dreaming and the instability of sleep: altered nonrapid eye movement sleep microstructure in individuals with frequent nightmares as revealed by the cyclic alternating pattern. *Sleep* 36, 413–419. doi: 10.5665/sleep.2462
- Sjöström, N., Hetta, J., and Waern, M. (2009). Persistent nightmares are associated with repeat suicide attempt. A prospective study. *Psychiatry Res.* 170, 208–221. doi: 10.1016/j.psychres.2008.09.006
- Solms, M. (1997). *The Neuropsychology of Dreams: A Clinico-anatomical Study*. Hillsdale, NJ: Erlbaum.
- Solms, M. (2000). Dreaming and REM sleep are controlled by different brain mechanisms. *Behav. Brain Sci.* 2, 843–850. doi: 10.1017/S0140525X00003988
- Solms, M. (2011). Neurobiology and the neurological basis of dreaming. *Handb. Clin. Neurol.* 98, 519–544. doi: 10.1016/B978-0-444-52006-7.00034-4
- Sopp, M. R., Michael, T., and Mecklinger, A. (2018). Effects of early morning nap sleep on associative memory for neutral and emotional stimuli. *Brain Res.* 1698, 29–42. doi: 10.1016/j.brainres.2018.06.020
- Sopp, M. R., Michael, T., Weeß, H. G., and Mecklinger, A. (2017). Remembering specific features of emotional events across time: the role of REM sleep and prefrontal theta oscillations. *Cogn. Affect. Behav. Neurosci.* 17, 1186–1209. doi: 10.3758/s13415-017-0542-8
- Spoormaker, V. I., Gleiser, P. M., and Cizisch, M. (2012). Frontoparietal connectivity and hierarchical structure of the brain’s functional network during sleep. *Front. Neurol.* 3:80. doi: 10.3389/fneur.2012.00080
- Spoormaker, V. I., Gvozdanovic, G. A., Sämann, P. G., and Cizisch, M. (2014). Ventromedial prefrontal cortex activity and rapid eye movement sleep are associated with subsequent fear expression in human subjects. *Exp. Brain Res.* 232, 1547–1554. doi: 10.1007/s00221-014-3831-2

- Spoormaker, V. I., and Montgomery, P. (2008). Disturbed sleep in post-traumatic stress disorder: secondary symptom or core feature? *Sleep Med. Rev.* 12, 169–184. doi: 10.1016/j.jpuro.2010.08.002
- Spoormaker, V. I., Schroter, M. S., Gleiser, P. M., Andrade, K. C., Dresler, M., Wehrle, R., et al. (2010). Development of a large-scale functional brain network during human non-rapid eye movement sleep. *J. Neurosci.* 30, 11379–11387. doi: 10.1523/JNEUROSCI.2015-10.2010
- Spoormaker, V. I., and Van den Bout, J. (2006). Lucid dreaming treatment for nightmares: a pilot study. *Psychother. Psychosom.* 75, 389–394. doi: 10.1159/000095446
- Spoormaker, V. I., Van den Bout, J., and Meijer, E. J. G. (2003). Lucid dreaming treatment for nightmares: a series of cases. *Dreaming* 13, 181–186. doi: 10.1023/A:1025325529560
- Sterpenich, V., Albouy, G., Boly, M., Vandewalle, G., Darsaud, A., Balet, E., et al. (2007). Sleep-related hippocampo-cortical interplay during emotional memory recollection. *PLoS Biol.* 5:e282. doi: 10.1371/journal.pbio.0050282
- Sterpenich, V., Perogamvros, L., Tononi, G., and Schwartz, S. (2019). Fear in dreams and in wakefulness: Evidence for day/night affective homeostasis. *bioRxiv*:534099. doi: 10.1101/534099
- Stickgold, R., Hobson, J. A., Fosse, R., and Fosse, M. (2001). Sleep, learning, and dreams: off-line memory reprocessing. *Science* 294, 1052–1057. doi: 10.1126/science.1063530
- Stumbrys, T., Erlacher, D., Schädlich, M., and Schredl, M. (2012). Induction of lucid dreams: a systematic review of evidence. *Conscious. Cogn.* 21, 1456–1475. doi: 10.1016/j.concog.2012.07.003
- Takeuchi, T., Ogilvie, R. D., Murphy, T. I., and Ferrelli, A. V. (2003). EEG activities during elicited sleep onset REM and NREM periods reflect different mechanisms of dream generation. *Clin. Neurophysiol.* 114, 210–220. doi: 10.1016/S1388-2457(02)00385-1
- Tamietto, M., and De Gelder, B. (2010). Neural bases of the non-conscious perception of emotional signals. *Nat. Rev. Neurosci.* 11, 697–709. doi: 10.1038/nrn2889
- Taub, J. M. (1971). The sleep-wakefulness cycle in Mexican adults. *J. Cross Cult. Psychol.* 2, 353–364. doi: 10.1177/002202217100200405
- Tort, A. B. L., Komorowski, R., Eichenbaum, H., and Kopell, N. (2010). Measuring phase-amplitude coupling between neuronal oscillations of different frequencies. *J. Neurophysiol.* 104, 1195–1210. doi: 10.1152/jn.00106.2010
- Tort, A. B. L., Komorowski, R. W., Manns, J. R., Kopell, N. J., and Eichenbaum, H. (2009). Theta-gamma coupling increases during the learning of item-context associations. *Proc. Natl. Acad. Sci. USA.* 106, 20942–20946. doi: 10.1073/pnas.0911331106
- Uribe, C. E., Garcia, A., and Tomaz, C. (2011). Electroencephalographic brain dynamics of memory encoding in emotionally arousing context. *Front. Behav. Neurosci.* 5:35. doi: 10.3389/fnbeh.2011.00035
- Vallat, R., Chatard, B., Blagrove, M., and Ruby, P. (2017). Characteristics of the memory sources of dreams: a new version of the content-matching paradigm to take mundane and remote memories into account. *PLoS ONE.* 12:e0185262. doi: 10.1371/journal.pone.0185262
- Vallat, R., Eichenlaub, J. B., Nicolas, A., and Ruby, P. (2018a). Dream recall frequency is associated with medial prefrontal cortex white-matter density. *Front. Psychol.* 9:1856. doi: 10.3389/fpsyg.2018.01856
- Vallat, R., Eskinazi, M., Nicolas, A., and Ruby, P. (2018b). Sleep and dream habits in a sample of French college students who report no sleep disorders. *J. Sleep Res.* 27:e12659. doi: 10.1111/jsr.12659
- van der Helm, E., Yao, J., Dutt, S., Rao, V., Saletin, J. M., and Walker, M. P. (2011). REM sleep depotentiates amygdala activity to previous emotional experiences. *Curr. Biol.* 21, 2029–2032. doi: 10.1016/j.cub.2011.10.052
- Van Overwalle, F., and Vandekerckhove, M. (2013). Implicit and explicit social mentalizing: dual processes driven by a shared neural network. *Front. Hum. Neurosci.* 7:560. doi: 10.3389/fnhum.2013.00560
- van Rijn, E., Eichenlaub, J. B., Lewis, P. A., Walker, M. P., Gaskell, M. G., Malinowski, J. E., et al. (2015). The dream-lag effect: selective processing of personally significant events during rapid eye movement sleep, but not during slow wave sleep. *Neurobiol. Learn. Mem.* 122, 98–109. doi: 10.1016/j.nlm.2015.01.009
- Vogel, G. W. (1976). Mentation reported from naps of narcoleptics. *Adv. Sleep Res.* 3, 161–168.
- Voss, U., Holzmann, R., Hobson, A., Paulus, W., Koppehele-Gossel, J., Klimke, A., et al. (2014). Induction of self-awareness in dreams through frontal low current stimulation of gamma activity. *Nat. Neurosci.* 17, 810–812. doi: 10.1038/nn.3719
- Voss, U., and Klimke, A. (2018). Dreaming during REM sleep: autobiographically meaningful or a simple reflection of a Hebbian-based memory consolidation process? *Arch. Ital. Biol.* 156, 99–111. doi: 10.1287/00039829201832
- Vuilleumier, P., Armony, J. L., Driver, J., and Dolan, R. J. (2001). Effects of attention and emotion on face processing in the human brain: an event-related fMRI study. *Neuron* 30, 829–841. doi: 10.1016/j.tics.2005.10.011
- Wagner, U., Gais, S., and Born, J. (2001). Emotional memory formation is enhanced across sleep intervals with high amounts of rapid eye movement sleep. *Learn. Memory* 8, 112–119. doi: 10.1101/lm.36801
- Walker, M. P. (2009). The role of sleep in cognition and emotion. *Ann. N. Y. Acad. Sci.* 1156, 168–197. doi: 10.1111/j.1749-6632.2009.04416.x
- Walker, M. P., and van der Helm, E. (2009). Overnight Therapy? The role of sleep in emotional brain processing. *Psychol. Bull.* 135, 731–748. doi: 10.1037/a0016570
- Wamsley, E. J., Donjacour, C. E., Scammell, T. E., Lammers, G. J., and Stickgold, R. (2014). Delusional confusion of dreaming and reality in narcolepsy. *Sleep* 37, 419–422. doi: 10.5665/sleep.3428
- Wamsley, E. J., and Stickgold, R. (2011). Memory, sleep, and dreaming: experiencing consolidation. *Sleep Med. Clin.* 6, 97–108. doi: 10.1016/j.jsmc.2010.12.008
- Wamsley, E. J., Tucker, M., Payne, J. D., Benavides, J. A., and Stickgold, R. (2010). Dreaming of a learning task is associated with enhanced sleep-dependent memory consolidation. *Curr. Biol.* 20, 850–855. doi: 10.1016/j.cub.2010.03.027
- White, T. P., Jansen, M., Doerge, K., Mullinger, K. J., Park, S. B., Liddle, E. B., et al. (2013). Theta power during encoding predicts subsequent-memory performance and default mode network deactivation. *Hum. Brain Mapp.* 34, 2929–2943. doi: 10.1002/hbm.22114
- Wurts, S. W., and Edgar, D. M. (2000). Circadian and homeostatic control of rapid eye movement (REM) sleep: promotion of REM tendency by the suprachiasmatic nucleus. *J. Neurosci.* 20, 4300–4310. doi: 10.1523/JNEUROSCI.20-11-04300.2000
- Ye, H., Chen, S., Huang, D., Zheng, H., Jia, Y., and Luo, J. (2015). Modulation of neural activity in the temporoparietal junction with transcranial direct current stimulation changes the role of beliefs in moral judgment. *Front. Hum. Neurosci.* 9:659. doi: 10.3389/fnhum.2015.00659
- Yetkin, S., Aydin, H., and Özgen, F. (2010). Polysomnography in patients with post-traumatic stress disorder. *Psychiatry Clin. Neurosci.* 64, 309–317. doi: 10.1111/j.1440-1819.2010.02084.x
- Yordanova, Y. N., Duffau, H., and Herbet, G. (2017). Neural pathways subserving face-based mentalizing. *Brain Struct. Funct.* 222, 3087–3105. doi: 10.1007/s00429-017-1388-0
- Young, C. K., and Eggermont, J. J. (2009). Coupling of mesoscopic brain oscillations: recent advances in analytical and theoretical perspectives. *Progr. Neurobiol.* 89, 61–78. doi: 10.1016/j.pneurobio.2009.06.002
- Young, L., Dodel-Feder, D., and Saxe, R. (2010). What gets the attention of the temporo-parietal junction? An fMRI investigation of attention and theory of mind. *Neuropsychologia* 48, 2658–2664. doi: 10.1016/j.neuropsychologia.2010.05.012
- Zadra, A. L., and Pihl, R. O. (1997). Lucid dreaming as a treatment for recurrent nightmares. *Psychother. Psychosom.* 66, 50–55. doi: 10.1159/000289106

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Scarpelli, Bartolacci, D'Atri, Gorgoni and De Gennaro. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.